



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> :C07D 239/50, 239/48, 239/46, 239/42,  
401/12, A61K 31/505

A1

(11) International Publication Number:

WO 99/50250

(43) International Publication Date:

7 October 1999 (07.10.99)

(21) International Application Number: PCT/EP99/02043

(22) International Filing Date: 24 March 1999 (24.03.99)

(30) Priority Data:

60/079,632	27 March 1998 (27.03.98)	US
98201587.7	14 May 1998 (14.05.98)	EP
98203948.9	25 November 1998 (25.11.98)	EP

(71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ANDRIES, Koenraad, Jozef, Lodewijk, Marcel [BE/BE]; Oosteneinde 9, B-2340 Beerse (BE). DE CORTE, Bart [BE/US]; 1590 Winding Road, Southampton, PA 18966 (US). DE JONGE, Marc, René [NL/NL]; Hontenissestraat 3, NL-5045 CA Tilburg (NL). HEERES, Jan [NL/BE]; Leemskuilen 18, B-2350 Vosselaar (BE). HO, Chih, Yung [US/US]; 600 Drinnon Way, Lansdale, PA 19446 (US). JANSSEN, Marcel, August, Constant [BE/BE]; Karel Druytslaan 15, B-2350 Vosselaar (BE). JANSSEN, Paul, Adriaan, Jan [BE/BE]; Antwerpses-teenweg 20, B-2350 Vosselaar (BE). KOYMANS, Lucien, Maria, Henricus [NL/BE]; Stationstraat 68 bus 4,

B-2300 Turnhout (BE). KUKLA, Michael, Joseph [US/US]; 1551 Oak Hollow Drive, Maple Glen, PA 19002 (US). LUDOVICI, Donald, William [US/US]; 1390 Masi Road, Quakertown, PA 18951 (US). VAN AKEN, Koen, Jeanne, Alfons [BE/BE]; Tuinbouwstraat 22, bus 2, B-2300 Turnhout (BE).

(74) Common Representative: JANSSEN PHARMACEUTICA N.V.; Att. Frank Daelemans, Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published


With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: HIV INHIBITING PYRIMIDINE DERIVATIVES

(57) Abstract

This invention concerns the use of the compounds of formula (I), the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein A is CH, CR<sup>4</sup> or N; n is 0 to 4; Q is hydrogen or -NR<sup>1</sup>R<sup>2</sup>; R<sup>1</sup> and R<sup>2</sup> are selected from hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)- amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl wherein each C<sub>1-12</sub>alkyl may optionally be substituted; or R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)aminoC<sub>1-4</sub>alkylidene; R<sup>3</sup> is hydrogen, aryl, C<sub>1-6</sub>alkylcarbonyl, optionally substituted C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl; and R<sup>4</sup> is hydroxy, halo, optionally substituted C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy; R<sup>5</sup> is hydrogen or C<sub>1-4</sub>alkyl; L is optionally substituted C<sub>1-10</sub>alkyl, C<sub>3-10</sub>alkenyl, C<sub>3-10</sub>alkynyl, C<sub>3-7</sub>cycloalkyl; or L is -X<sup>1</sup>R<sup>6</sup> or -X<sup>2</sup>-Alk-R<sup>7</sup> wherein R<sup>6</sup> and R<sup>7</sup> are optionally substituted phenyl; X<sup>1</sup> and X<sup>2</sup> are -NR<sup>3</sup>-, -NH-NH-, -N=N-, -O-, -S-, -S(=O)- or -S(=O)<sub>2</sub>-; Alk is C<sub>1-4</sub>alkanediyl; aryl is optionally substituted phenyl; Het is an optionally substituted aliphatic or aromatic heterocyclic radical; for the manufacture of a medicine for the treatment of subjects suffering from HIV (Human Immunodeficiency Virus) infection. It further relates to new compounds being a subgroup of the compounds of formula (I), their preparation and compositions comprising them.



(1)

-1-

## HIV INHIBITING PYRIMIDINE DERIVATIVES

5 The present invention is concerned with pyrimidine derivatives having HIV replication inhibiting properties. The invention further relates to methods for their preparation and pharmaceutical compositions comprising them. The invention also relates to the use of said compounds in the manufacture of a medicament useful for the treatment of subjects suffering from HIV (Human Immunodeficiency Virus) infection.

10 Compounds structurally related to the present compounds are disclosed in the prior art.


JP-2,052,360, JP-2,308,248, JP-9,080,676 and JP-9,068,784 disclose a number of trisubstituted pyrimidines useful in photographic material. JP-8,199,163 discloses trisubstituted pyrimidines useful in an organic electroluminescent device. JP-2,300,264 and GB-1,477,349 disclose pyrimidinetriamines for their use in the dye industry.

J. Indian Chem. Soc. (1975), 52(8), 774-775 discloses the synthesis of some bis(arylamino)pyrimidines. J. Heterocycl. Chem. (1973), 10(2), 167-171 discloses the condensation of various aminopyrimidines with picryl halides. J. Org. Chem. (1961), 26, 4433-4440 discloses several triaminopyrimidines as intermediates in the synthesis of triazolo[4,5-d]pyrimidines.

WO 91/18887 discloses diaminopyrimidines as gastric acid secretion inhibitors.

25 Unexpectedly, it has now been found that the compounds of formula (I) effectively inhibit the replication of the Human Immunodeficiency Virus (HIV) and consequently may be useful for the treatment of individuals infected by HIV.

The present invention concerns the use of the compounds of formula



30

the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

A is CH, CR<sup>4</sup> or N;

n is 0, 1, 2, 3 or 4;

35 Q is hydrogen or -NR<sup>1</sup>R<sup>2</sup>;

$R^1$  and  $R^2$  are each independently selected from hydrogen, hydroxy,  $C_{1-12}$ alkyl,  $C_{1-12}$ alkyloxy,  $C_{1-12}$ alkylcarbonyl,  $C_{1-12}$ alkyloxycarbonyl, aryl, amino, mono- or di( $C_{1-12}$ alkyl)amino, mono- or di( $C_{1-12}$ alkyl)aminocarbonyl wherein each of the  
 5       aforementioned  $C_{1-12}$ alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy,  $C_{1-6}$ alkyloxy, hydroxy $C_{1-6}$ alkyloxy, carboxyl,  $C_{1-6}$ alkyloxycarbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonylamino, mono- or di( $C_{1-6}$ alkyl)amino, aryl and Het; or

$R^1$  and  $R^2$  taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or  
 10       mono- or di( $C_{1-12}$ alkyl)amino $C_{1-4}$ alkylidene;

$R^3$  is hydrogen, aryl,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkyl substituted with  $C_{1-6}$ alkyloxycarbonyl; and

each  $R^4$  independently is hydroxy, halo,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, amino-carbonyl, nitro, amino, trihalomethyl, trihalomethyloxy or  $C_{1-6}$ alkyl substituted with  
 15       cyano or aminocarbonyl;

$R^5$  is hydrogen or  $C_{1-4}$ alkyl;

L is  $C_{1-10}$ alkyl,  $C_{3-10}$ alkenyl,  $C_{3-10}$ alkynyl,  $C_{3-7}$ cycloalkyl, or  $C_{1-10}$ alkyl substituted with one or two substituents independently selected from  $C_{3-7}$ cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two,  
 20       three, four or where possible five substituents each independently selected from halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, aminocarbonyl,  $C_{1-6}$ alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl, trihalomethyloxy and  $C_{1-6}$ alkylcarbonyl; or

L is  $-X^1-R^6$  or  $-X^2-Alk-R^7$  wherein

$R^6$  and  $R^7$  each independently are phenyl or phenyl substituted with one, two, three,  
 25       four or five substituents each independently selected from halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyloxycarbonyl, formyl, cyano, aminocarbonyl, nitro, amino, trihalomethyloxy and trihalomethyl; and

$X^1$  and  $X^2$  are each independently  $-NR^3-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-S-$ ,  $-S(=O)-$  or  $-S(=O)_2-$ ;

30       Alk is  $C_{1-4}$ alkanediyl;

aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, nitro and trifluoromethyl;


Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is  
 35       selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic hetero-

cyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy;

for the manufacture of a medicine for the treatment of subjects suffering from HIV  
5 (Human Immunodeficiency Virus) infection.

The present invention also relates to a method of treating warm-blooded animals suffering from HIV (Human Immunodeficiency Virus) infection. Said method comprises the administration of a therapeutically effective amount of a compound of  
10 formula (I) or any subgroup thereof, a *N*-oxide form, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof in admixture with a pharmaceutical carrier.

This invention also concerns compounds of formula



15

the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein L, Q, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and A are as defined under formula (I), and

R<sup>4'</sup> is halo, C<sub>1-6</sub>alkyl, cyano, aminocarbonyl, nitro, trihalomethyl, trihalomethyloxy or  
20 C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;


n' is 0, 1, 2 or 3;

with the proviso that Q and L are other than anilino, 2,4,6-trinitro-anilino, 3-methoxy-anilino, 4-methoxy-anilino, 3,4-dimethoxy-anilino, 3-chloro-4-fluoro-anilino, 4-cyano-anilino, 2-(C<sub>1-6</sub>alkyl)-anilino, 4-(C<sub>1-6</sub>alkyl)-anilino, 3-chloro-anilino, 4-bromo-anilino,  
25 4-nitro-anilino and 4-chloro-anilino.

As used in the foregoing definitions and hereinafter halo defines fluoro, chloro, bromo and iodo; C<sub>1-4</sub>alkyl as a group or part of a group encompasses the straight and branched chained saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for  
30 example, methyl, ethyl, propyl, butyl and the like; C<sub>1-6</sub>alkyl as a group or part of a group encompasses the straight and branched chained saturated hydrocarbon radicals as defined in C<sub>1-4</sub>alkyl as well as the higher homologues thereof containing 5 or 6 carbon atoms such as, for example pentyl or hexyl; C<sub>1-10</sub>alkyl as a group or part of a group


group encompasses the straight and branched chained saturated hydrocarbon radicals as defined in C<sub>1-6</sub>alkyl as well as the higher homologues thereof containing 7 to 10 carbon atoms such as, for example, heptyl, octyl, nonyl or decyl; C<sub>1-12</sub>alkyl as a group or part of a group encompasses the straight and branched chained saturated hydrocarbon radicals as defined in C<sub>1-10</sub>alkyl as well as the higher homologues thereof containing 11 or 12 carbon atoms such as, for example, undecyl, dodecyl and the like; C<sub>1-4</sub>alkylidene as a group or part of a group defines bivalent straight and branched chained hydrocarbons having from 1 to 4 carbon atoms such as, for example, methylene, ethylidene, propylidene, butylidene and the like; C<sub>1-4</sub>alkanediyl as a group or part of a group encompasses those radicals defined under C<sub>1-4</sub>alkylidene as well as other bivalent straight and branched chained hydrocarbons having from 1 to 4 carbon atoms such as, for example, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl and the like; C<sub>3-7</sub>cycloalkyl as a group or part of a group is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; C<sub>3-10</sub>alkenyl as a group or part of a group defines straight and branch chained hydrocarbon radicals containing one double bond and having from 3 to 10 carbon atoms such as, for example, 2-propenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl, 3-heptenyl, 2-octenyl, 2-nonenyl, 2-decenyl and the like, whereby the carbon atom attached to the pyrimidine ring is preferably an aliphatic carbon atom; C<sub>3-10</sub>alkynyl as a group or part of a group defines straight and branch chained hydrocarbon radicals containing one triple bond and having from 3 to 10 carbon atoms such as, for example, 2-propynyl, 2-butylnyl, 2-pentylnyl, 3-pentylnyl, 3-methyl-2-butylnyl, 3-hexynyl, 3-heptynyl, 2-octynyl, 2-nonylnyl, 2-decynyl and the like, whereby the carbon atom attached to the pyrimidine ring is preferably an aliphatic carbon atom.

It is to be understood that the three substituents [L, Q and NR<sup>3</sup>(optionally substituted phenyl or pyridyl)] on the pyrimidine ring can be on any free position of the pyrimidine ring. Thus, given the following numbering of the pyrimidine ring




the three substituents may be connected to the pyrimidine ring in three different ways :  
 2-L, 4-Q, 6-NR<sup>3</sup>(optionally substituted phenyl or pyridyl); or  
 4-L, 2-Q, 6-NR<sup>3</sup>(optionally substituted phenyl or pyridyl); or  
 6-L, 4-Q, 2-NR<sup>3</sup>(optionally substituted phenyl or pyridyl).

The positions 4 and 6 are equivalent to one another. For instance, substitution pattern 6-L, 4-Q, 2-NR<sup>3</sup>(optionally substituted phenyl or pyridyl), which is a preferred substitution pattern, is equivalent to substitution pattern 4-L, 6-Q, 2-NR<sup>3</sup>(optionally substituted phenyl or pyridyl). Said subgroup of compounds is represented by formula



5

An interesting group of compounds are the compounds of formula



10

Of particular interest are those compounds of formula (I'-1) wherein L and Q are other than anilino, 2,4,6-trinitro-anilino, 4-(C<sub>1-6</sub>alkyl)-anilino, 4-bromo-anilino, 4-nitro-anilino and 4-chloro-anilino; and of more particular interest are those compounds of formula (I'-1) wherein R<sup>4'</sup> is cyano, aminocarbonyl or C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl.

15

The addition salts as mentioned herein are meant to comprise the therapeutically active addition salt forms which the compounds of the present invention are able to form with appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-amino-salicylic, pamoic and the like acids.

20

The pharmaceutically acceptable addition salts as mentioned hereinabove are also meant to comprise the therapeutically active non-toxic base, in particular, a metal or amine addition salt forms which the compounds of the present invention are able to form. Said salts can conveniently be obtained by treating the compounds of the present invention containing acidic hydrogen atoms with appropriate organic and inorganic bases such as, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts

25

with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

Conversely said salt forms can be converted by treatment with an appropriate base or acid into the free acid or base form.

5

The term addition salts also comprises the hydrates and the solvent addition forms which the compounds of the present invention are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

10 The term stereochemically isomeric forms of compounds of the present invention, as used hereinbefore, defines all possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of the present invention may possess.

Unless otherwise mentioned or indicated, the chemical designation of a compound  
15 encompasses the mixture of all possible stereochemically isomeric forms which said compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of the present invention both in pure form or in admixture with each other are intended to be embraced within the scope of the present  
20 invention.

Some of the compounds of the present invention may also exist in their tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

25

Whenever used hereinafter, the term "compounds of the present invention" is meant to include the compounds of formula (I), (I-1), (I'), (I'-1) or any subgroup thereof, also the *N*-oxides, the pharmaceutically acceptable addition salts and all stereoisomeric forms.

30 The group containing those compounds of the present invention wherein Q is  $NR^1R^2$ , each  $R^4$  independently is hydroxy, halo,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, amino-carbonyl, nitro, amino, trihalomethyl or trihalomethyloxy ; L is  $C_{1-10}$ alkyl,  $C_{3-10}$ alkenyl,  $C_{3-10}$ alkynyl,  $C_{3-7}$ cycloalkyl, or  $C_{1-10}$ alkyl substituted with one or two substituents independently selected from  $C_{3-7}$ cycloalkyl, indolyl or indolyl substituted with one,  
35 two, three or four substituents each independently selected from halo,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy and  $C_{1-6}$ alkylcarbonyl, phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy,

cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy and C<sub>1-6</sub>alkyl-carbonyl; or L is -X<sup>1</sup>-R<sup>6</sup> wherein R<sup>6</sup> is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylcarbonyl, cyano, nitro and trifluoromethyl; is of interest.

5

Also of interest is the group containing those compounds of the present invention wherein Q is NR<sup>1</sup>R<sup>2</sup>; each R<sup>4</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethoxy; L is C<sub>1-10</sub>alkyl substituted with one or two substituents independently selected from phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy and C<sub>1-6</sub>alkylcarbonyl; or L is -X<sup>1</sup>-R<sup>6</sup> wherein R<sup>6</sup> is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylcarbonyl, cyano, nitro and trifluoromethyl; with the proviso that compounds

10

15

20

25

30

35

- (a) *N*2-hydroxy-*N*2-methyl-*N*4,*N*6-diphenyl-2,4,6-pyrimidinetriamine;
- (b) *N,N,N',N',N'',N'''*-hexakis(3-methylphenyl)-2,4,6-pyrimidinetriamine;
- (c) *N*4-methyl-*N*2-(2-methylphenyl)-*N*4-phenyl-2,4,6-pyrimidinetriamine;
- (d) *N*4-methyl-*N*2-(2-methylphenyl)-*N*4-phenyl-6-(phenylmethyl)-2,4-pyrimidine diamine;
- (e) *N*4-(2-methylphenyl)-6-(phenylmethyl)-2,4-pyrimidinediamine;
- (f) *N,N,N'''*-tris(4-methoxyphenyl)-2,4,6-pyrimidinetriamine;
- (g) *N,N'*-bis(4-hexylphenyl)-6-(4-methoxyphenoxy)-2,4-pyrimidinediamine;
- (h) *N*2,*N*4-bis(4-hexylphenyl)-*N*6,*N*6-dimethyl-2,4,6-pyrimidinetriamine;
- (i) *N,N,N'''*-tris(4-hexylphenyl)-2,4,6-pyrimidinetriamine;
- (j) *N*2,*N*2-dimethyl-*N*4,*N*6-bis(4-methylphenyl)-2,4,6-pyrimidinetriamine;
- (k) *N,N,N'*-tris(4-methylphenyl)-2,4,6-pyrimidinetriamine;
- (l) *N,N,N'*-triphenyl-2,4,6-pyrimidinetriamine;
- (m) *N,N,N',N',N'',N'''*-hexakis(4-ethoxyphenyl)-2,4,6-pyrimidinetriamine;
- (n) *N*4,*N*6-bis(2-chlorophenyl)-2,4,6-pyrimidinetriamine;
- (o) *N*4,*N*6-bis(3-chlorophenyl)-2,4,6-pyrimidinetriamine;
- (p) *N*4,*N*6-bis(2-ethoxyphenyl)-2,4,6-pyrimidinetriamine;
- (q) *N*4,*N*6-bis(4-ethoxyphenyl)-2,4,6-pyrimidinetriamine;
- (r) *N*4,*N*6-bis(2-methylphenyl)-2,4,6-pyrimidinetriamine;
- (s) *N*4,*N*6-bis(4-bromophenyl)-2,4,6-pyrimidinetriamine;
- (t) *N*4,*N*6-bis(4-methylphenyl)-2,4,6-pyrimidinetriamine;
- (u) *N*2,*N*4-bis(4-methoxyphenyl)-2,4,6-pyrimidinetriamine;

- (v) *N2,N4*-bis(4-methylphenyl)-2,4,6-pyrimidinetriamine;
- (w) *N,N,N'*-tris(2,4,6-trinitrophenyl)-2,4,6-pyrimidinetriamine;
- (x) *N4,N6*-bis(4-chlorophenyl)-2,4,6-pyrimidinetriamine;
- (y) *N4,N6*-bis(4-methoxyphenyl)-2,4,6-pyrimidinetriamine;
- 5 (z) *N2,N4,N6*-trimethyl-*N2,N4,N6*-triphenyl pyrimidine-2,4,6-triyltriamine;
- (aa) *N4,N4*-dimethyl-*N2,N6*-di-*p*-tolyl-pyrimidine-2,4,6-triyltriamine; and
- (bb) *N2,N4*-diphenyl-pyrimidine-2,4,6-triyltriamine are not included.

10 Suitably, Q may also be hydrogen in the above two groups of interest.

A special group of compounds are those compounds of formula (I) or (I') wherein n is at least 1 and at least one R<sup>4</sup> is cyano; preferably, n is 1 and R<sup>4</sup> is cyano substituted in the para position relative to the NR<sup>3</sup> moiety.

15 Another special group of compounds contains those compounds of formula (I) or (I') which are other than

- (c) *N4*-methyl-*N2*-(2-methylphenyl)-*N4*-phenyl-2,4,6-pyrimidinetriamine;
- (d) *N4*-methyl-*N2*-(2-methylphenyl)-*N4*-phenyl-6-(phenylmethyl)-2,4-pyrimidine-diamine;
- 20 (e) *N4*-(2-methylphenyl)-6-(phenylmethyl)-2,4-pyrimidinediamine; the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof.

25 An interesting group of compounds are those compounds of the present invention wherein the NR<sup>3</sup>(substituted phenyl or pyridyl) moiety is in the 4- or 6-position of the pyrimidine ring.

30 Another interesting group are those compounds of the present invention wherein each R<sup>4</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethyloxy; R<sup>6</sup> is phenyl or phenyl substituted with one, two or three, four or five substituents each independently selected from halo, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylcarbonyl, cyano, nitro and trifluoromethyl; and aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C<sub>1-6</sub>alkyloxy, cyano, nitro and trifluoromethyl.

35 Suitably, Q is NR<sup>1</sup>R<sup>2</sup> wherein R<sup>1</sup> is hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl; and R<sup>2</sup> is hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)amino,

mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl; wherein each of the aforementioned C<sub>1-12</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonyl-amino, mono- or di(C<sub>1-6</sub>alkyl)amino, aryl and Het; or R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)amino-C<sub>1-4</sub>alkylidene.

Suitably, L is C<sub>1-10</sub>alkyl substituted with one or two substituents independently selected from C<sub>3-7</sub>cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two, three, four or where possible five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, C<sub>1-6</sub>alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl, trihalomethoxy and C<sub>1-6</sub>alkylcarbonyl; or L is -X<sup>1</sup>-R<sup>6</sup> or -X<sup>2</sup>-Alk-R<sup>7</sup> and when X<sup>1</sup> is NR<sup>3</sup>, then R<sup>6</sup> is phenyl substituted with one, two, three, four or five substituents each independently selected from C<sub>1-6</sub>alkyloxycarbonyl, formyl, nitro and trihalomethoxy.

Suitably, R<sup>4</sup> or R<sup>4'</sup> is nitro, trihalomethoxy or C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl.

Suitably, R<sup>6</sup> is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, formyl, cyano, aminocarbonyl, nitro, amino, trihalomethoxy and trihalomethyl.

Suitably, both Q and R<sup>5</sup> are hydrogen.

Suitably, L is C<sub>1-10</sub>alkyl substituted with one or two substituents independently selected from C<sub>3-7</sub>cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two, three, four or where possible five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, C<sub>1-6</sub>alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl, trihalomethoxy and C<sub>1-6</sub>alkylcarbonyl; or L is -X<sup>1</sup>-R<sup>6</sup> or -X<sup>2</sup>-Alk-R<sup>7</sup>; and R<sup>5</sup> is hydrogen.

Particular groups of compounds are those groups wherein one or more of the following conditions are met :

- (i) n is 0, 1, 2 or 3;
- (ii) Q is hydrogen;

- (iii) Q is  $\text{NR}^1\text{R}^2$  wherein  $\text{R}^1$  and  $\text{R}^2$  are each independently selected from hydrogen, hydroxy,  $\text{C}_{1-12}$ alkyl,  $\text{C}_{1-12}$ alkyloxy,  $\text{C}_{1-12}$ alkylcarbonyl,  $\text{C}_{1-12}$ alkyloxycarbonyl, cyano wherein the aforementioned  $\text{C}_{1-12}$ alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, cyano,  $\text{C}_{1-6}$ alkyloxy, hydroxy $\text{C}_{1-6}$ alkyloxy, aryl and Het; or  $\text{R}^1$  and  $\text{R}^2$  taken together may form mono- or di( $\text{C}_{1-12}$ alkyl)amino $\text{C}_{1-4}$ alkylidene;
- (iv)  $\text{R}^3$  is hydrogen or  $\text{C}_{1-6}$ alkyl;
- (v)  $\text{R}^4$  is cyano, aminocarbonyl, amino, nitro, hydroxy, halo,  $\text{C}_{1-6}$ alkyl or cyano $\text{C}_{1-6}$ alkyl;
- (vi)  $\text{R}^4$  is cyano, aminocarbonyl, halo,  $\text{C}_{1-6}$ alkyl or cyano $\text{C}_{1-6}$ alkyl;
- (vii)  $\text{R}^5$  is hydrogen or methyl;
- (viii) L is  $\text{C}_{1-10}$ alkyl substituted with phenyl substituted with one or two halogens; or L is  $-\text{X}^1-\text{R}^6$  wherein  $\text{R}^6$  is phenyl substituted with one, two or three substituents selected from  $\text{C}_{1-6}$ alkyl, trifluoromethyl, trifluoromethoxy, cyano; and halogen, and  $\text{X}^1$  is  $-\text{S}-$ ,  $-\text{O}-$  or  $-\text{NR}^3-$ ; or L is  $-\text{X}^2-\text{Alk}-\text{R}^7$  wherein  $\text{R}^7$  is phenyl substituted with one, two or three substituents selected from  $\text{C}_{1-6}$ alkyl, cyano, and halogen and  $\text{X}^2$  is NH.

Other particular compounds are those compounds of the present invention wherein L contains phenyl, 2,6-disubstituted-phenyl, 2,4,6-trisubstituted-phenyl or 2,3,4,5-tetra-substituted-phenyl; especially, L contains phenyl, 2,4,6-trihalo-phenyl, 2,4,6-tri $\text{C}_{1-4}$ alkyl-phenyl, 2,3,4,5-tetrahalo-phenyl, 2,4-dihalo-6- $\text{C}_{1-4}$ alkyl-phenyl, 2,6-dihalo-4- $\text{C}_{1-4}$ alkyl-phenyl, 2,6-dihalo-4-cyano-phenyl, 2,6-dihalo-4-trifluoromethoxy-phenyl, 2,6-dihalo-4-trifluoromethyl-phenyl, 2,6-di $\text{C}_{1-4}$ alkyl-4-halo-phenyl, 2,6-di $\text{C}_{1-4}$ alkyl-4-cyano-phenyl, 2,6-dihalo-phenyl or 2,6-di $\text{C}_{1-4}$ alkyl-phenyl; more in particular, L contains phenyl, 2,4,6-trichloro-phenyl, 2,4,6-trimethyl-phenyl, 2,4-dibromo-3,5-dichloro-phenyl, 2,4-dibromo-6-fluoro-phenyl, 2,4-dichloro-6-methyl-phenyl, 2,6-dibromo-4-isopropyl-phenyl, 2,6-dibromo-4-methyl-phenyl, 2,6-dibromo-4-prop-1-yl-phenyl, 2,6-dichloro-4-cyano-phenyl, 2,6-dichloro-4-trifluoromethoxy-phenyl, 2,6-dichloro-4-trifluoromethyl-phenyl, 2,6-dichloro-phenyl, 2,6-dimethyl-4-(1,1-dimethylethyl)-phenyl, 2,6-dimethyl-phenyl, 2-bromo-4-fluoro-6-methyl-phenyl, 2-bromo-6-chloro-4-fluoro-phenyl, 4-bromo-2,6-dimethyl-phenyl, 4-chloro-2,6-dimethyl-phenyl, or 4-cyano-2,6-dimethyl-phenyl.

More particular compounds are the compounds of the present invention wherein L is 2,6-dichlorobenzyl, or L is  $-\text{X}^1-\text{R}^6$  wherein  $\text{X}^1$  is  $-\text{NR}^3-$ ,  $-\text{S}-$  or  $-\text{O}-$  and  $\text{R}^6$  is 2,4,6-tri-

chlorophenyl, 2,4,6-trimethyl-phenyl, 2,4-dibromo-3,5-dichloro-phenyl, 2,4-dibromo-6-fluoro-phenyl, 2,4-dichloro-6-methyl-phenyl, 2,6-dibromo-4-isopropyl-phenyl, 2,6-dibromo-4-methyl-phenyl, 2,6-dibromo-4-prop-1-yl-phenyl, 2,6-dichloro-4-cyano-phenyl, 2,6-dichloro-4-trifluoromethoxy-phenyl, 2,6-dichloro-4-trifluoromethyl-phenyl, 2,6-dichloro-phenyl, 2,6-dimethyl-4-(1,1-dimethylethyl)-phenyl, 2,6-dimethyl-phenyl, 2-bromo-4-fluoro-6-methyl-phenyl, 2-bromo-6-chloro-4-fluoro-phenyl, 4-bromo-2,6-dimethyl-phenyl, 4-chloro-2,6-dimethyl-phenyl, 4-cyano-2,6-dimethyl-phenyl; or L is  $-X^2\text{-Alk-R}^7$  wherein  $-X^2\text{-Alk-}$  is  $-\text{NH-CH}_2\text{-}$  and  $\text{R}^7$  phenyl.

10 Still other particular compounds are those compounds of formula (I) where  $\text{R}^3$  is hydrogen, A is CH, n is 1, and  $\text{R}^4$  is halo, methyl or cyano and is positioned in the 4 position of the phenyl ring.

Preferred compounds are those compounds of the present invention wherein L is 2,6-dichlorobenzyl and the  $\text{NR}^3$  (optionally substituted phenyl or pyridyl) moiety represents *p*-cyano-anilino and is in the 2 position of the pyrimidine ring.

Other preferred compounds are those compounds of the present invention wherein Q is hydrogen, L is  $-X^1\text{-R}^6$  wherein  $\text{X}^1$  is  $-\text{NH-}$  and  $\text{R}^6$  is 2,4,6-trimethyl-phenyl or 4-cyano-2,6-dimethylphenyl, the  $\text{NR}^3$  (optionally substituted phenyl or pyridyl) moiety represents *p*-cyano-anilino and is in the 2 position of the pyrimidine ring.

Still other preferred compounds are those compounds of the present invention wherein L is  $-X^2\text{-Alk-R}^7$  wherein  $\text{X}^2$  is  $-\text{NH-}$ , Alk is methylene and  $\text{R}^7$  is phenyl, 2,6-dichloro-phenyl, 2,4,6-trimethyl-phenyl or 4-cyano-2,6-dimethylphenyl.

More preferred are those compounds of formula (I'-1) wherein  $\text{R}^{4'}$  is halo, cyano, aminocarbonyl or cyano $\text{C}_{1-6}$ alkyl; n is zero, A is CH,  $\text{R}^3$  is hydrogen;  $\text{R}^5$  is hydrogen or methyl; Q is hydrogen or  $\text{NHR}^1$ ; and L contains phenyl, 2,4,6-trichloro-phenyl, 2,4,6-trimethyl-phenyl, 2,4-dibromo-3,5-dichloro-phenyl, 2,4-dibromo-6-fluoro-phenyl, 2,4-dichloro-6-methyl-phenyl, 2,6-dibromo-4-isopropyl-phenyl, 2,6-dibromo-4-methyl-phenyl, 2,6-dibromo-4-prop-1-yl-phenyl, 2,6-dichloro-4-cyano-phenyl, 2,6-dichloro-4-trifluoromethoxy-phenyl, 2,6-dichloro-4-trifluoromethyl-phenyl, 2,6-dichloro-phenyl, 2,6-dimethyl-4-(1,1-dimethylethyl)-phenyl, 2,6-dimethyl-phenyl, 2-bromo-4-fluoro-6-methyl-phenyl, 2-bromo-6-chloro-4-fluoro-phenyl, 4-bromo-2,6-dimethyl-phenyl, 4-chloro-2,6-dimethyl-phenyl, or 4-cyano-2,6-dimethyl-phenyl.

Most preferred are

4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]benzonitrile;

- 6-[(2,6-dichlorophenyl)methyl]-*N*2-(4-fluorophenyl)-2,4-pyrimidinediamine;  
4-[[4-[(2,4-dichlorophenyl)methyl]-6-[(4-hydroxybutyl)amino]-2-pyrimidinyl]amino]-  
benzonitrile;  
4-[[4-[(2,6-dichlorophenyl)methyl]-6-[(3-hydroxypropyl)amino]-2-pyrimidinyl]amino]-  
5 benzonitrile;  
*N*-[2-[(4-cyanophenyl)amino]-6-[(2,6-dichlorophenyl)methyl]-4-pyrimidinyl]-  
acetamide;  
*N*-[2-[(4-cyanophenyl)amino]-6-[(2,6-dichlorophenyl)methyl]-4-pyrimidinyl]-  
butanamide;  
10 4-[[2-amino-6-(2,6-dichlorophenoxy)-4-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,6-dichlorophenyl)methyl]-6-[(2-hydroxy-2-phenylethyl)amino]-2-  
pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,6-dichlorophenyl)methyl]-6-[[3-(2-oxo-1-pyrrolidinyl)propyl]amino]-2-  
pyrimidinyl]amino]benzonitrile;  
15 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[[2-(2-hydroxyethoxy)ethyl]amino]-2-  
pyrimidinyl]amino]benzonitrile monohydrochloride;  
4-[[4-[(2,6-dichlorophenyl)methyl]-6-[(2,3-dihydroxypropyl)amino]-2-pyrimidinyl]-  
amino]benzonitrile;  
4-[[4-[(2,6-dichlorophenyl)methyl]-6-(hydroxyamino)-2-pyrimidinyl]amino]-  
20 benzonitrile;  
4-[[4-[(2-cyanoethyl)amino]-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]-  
benzonitrile;  
4-[[4-[(2,6-dichlorophenyl)methyl]-6-[[2-(1-pyrrolidinyl)ethyl]amino]-2-pyrimidinyl]-  
amino]benzonitrile;  
25 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-5-methyl-2-pyrimidinyl]amino]-  
benzonitrile;  
*N*2-(4-bromophenyl)-6-[(2,6-dichlorophenyl)methyl]-5-methyl-2,4-pyrimidinediamine;  
4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[2-[(2,4,6-trimethylphenyl)amino]-4-pyrimidinyl]amino]benzonitrile;  
30 4-[[4-[(2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-(2,4,6-trimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,6-dichlorophenyl)thio]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[[2,6-dibromo-4-(1-methylethyl)phenyl]amino]-2-pyrimidinyl]amino]-  
benzonitrile;  
35 4-[[4-[[2,6-dichloro-4-(trifluoromethyl)phenyl]amino]-2-pyrimidinyl]amino]-  
benzonitrile;  
4-[[4-[(2,4-dichloro-6-methylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

- 4-[[2-[(cyanophenyl)amino]-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;  
4-[[4-[(2,4-dibromo-6-fluorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-5-methyl-2-pyrimidinyl]amino]-  
benzeneacetonitrile;
- 5 4-[[4-[methyl(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,4,6-trichlorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,4,6-trimethylphenyl)thio]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-amino-6-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 10 4-[[2-amino-6-[(2,4,6-trimethylphenyl)amino]-4-pyrimidinyl]amino]benzonitrile;  
4-[[4-(2-bromo-4-chloro-6-methylphenoxy)-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(4-chloro-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
3,5-dichloro-4-[[2-[(4-cyanophenyl)amino]-4-pyrimidinyl]amino]benzonitrile;  
4-[[4-[[2,6-dichloro-4-(trifluoromethoxy)phenyl]amino]-2-pyrimidinyl]amino]-
- 15 benzonitrile;  
4-[[4-[(2,4-dibromo-3,6-dichlorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,6-dibromo-4-propylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzamide;  
4-[[4-[(4-(1,1-dimethylethyl)-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]-
- 20 benzonitrile;  
4-[[2-[(4-cyanophenyl)amino]-4-pyrimidinyl]oxy]-3,5-dimethylbenzonitrile;  
4-[[4-[(4-chloro-2,6-dimethylphenyl)amino]-5-methyl-2-pyrimidinyl]amino]-  
benzonitrile;  
4-[[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethyl-
- 25 benzonitrile;  
4-[[4-[[4-(1,1-dimethylethyl)-2,6-dimethylphenyl]amino]-5-methyl-2-pyrimidinyl]-  
amino]benzonitrile;  
4-[[4-[(4-bromo-2,6-dimethylphenyl)amino]-5-methyl-2-pyrimidinyl]amino]-  
benzonitrile;
- 30 4-[[5-methyl-4-[(2,4,6-trimethylphenyl)thio]-2-pyrimidinyl]amino]benzonitrile;  
4-[[4-[(2,6-dibromo-4-propylphenyl)amino]-5-methyl-2-pyrimidinyl]amino]-  
benzonitrile;  
4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzamide, N3-oxide;  
N2-(4-chlorophenyl)-N4-(2,4,6-trimethylphenyl)-2,4-pyrimidinediamine;
- 35 4-[[4-[[2,6-dibromo-4-(1-methylethyl)phenyl]amino]-5-methyl-2-pyrimidinyl]amino]-  
benzonitrile;  
4-[[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethyl

benzonitrile;


4-[[4-[(phenylmethyl)amino]-2-pyrimidinyl]amino]benzonitrile;

the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof.

5

The compounds of formula (I) can be prepared according to art-known procedures.


In particular, compounds of formula (I') can generally be prepared by reacting an intermediate of formula (II-A) wherein  $W^1$  is a suitable leaving group such as, for example, a halogen, with an amino derivative of formula (III) optionally in a solvent such as, for example, water, 2-propanol, diethylether, 1-methyl-2-pyrrolidinone and the like, and optionally in the presence of an acid such as, for example, 1 N hydrochloric acid in diethylether. It may be convenient to perform the reaction under a reaction-inert atmosphere such as, for example, oxygen free argon or nitrogen.



In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, distillation, trituration and chromatography.

20

Analogously to the reaction procedure described above,  $H-NR^1R^2$  (VI) can also be reacted with an intermediate of formula (II-B).




Suitable solvents for the above reaction include, for instance, 2-propanol or 1,4-dioxane.

In case Q is  $NR^1R^2$  and  $R^2$  contains a hydroxy moiety, it may be convenient to perform

the above reaction with a protected form of intermediate (VI) whereby the hydroxy moiety bears a suitable protecting group P being, for instance, a benzyl, and subsequently removing the protective group according to art-known methodologies, such as, for example, reacting with  $\text{BBr}_3$  in dichloromethane under nitrogen atmosphere.


It is also possible to react  $\text{H-X}^1\text{-R}^6$  with an intermediate of formula (II-C) in a suitable solvent such as, for example, 1,4-dioxane, thus obtaining compounds of formula (I') wherein L is  $-\text{X}^1\text{-R}^6$ , said compounds being represented by formula (I'-c).



Depending on the nature of  $\text{X}^1$  a suitable base or acid may be used to improve the reaction rate. For instance, in case  $\text{X}^1$  is  $-\text{O}-$ , sodium hydride may be used as suitable base; or in case  $\text{X}^1$  is  $\text{NR}^3$ ,  $\text{HCl}$  may be used as a suitable acid.

15 The compounds of formula (I') may further be prepared by converting compounds of formula (I') into each other according to art-known group transformation reactions.

For instance, compounds of formula (I') whereby Q is  $\text{NR}^1\text{R}^2$  and  $\text{R}^1$  and  $\text{R}^2$  are taken together to form mono- or di( $\text{C}_{1-12}$ alkyl)amino $\text{C}_{1-4}$ alkylidene, said compounds being represented by formula (I'-a), may be prepared by reacting a compound of formula (I') wherein  $\text{R}^1$  and  $\text{R}^2$  are hydrogen, with an intermediate of formula (IV) or a functional derivative thereof.



Also, compounds of formula (I') wherein Q is  $\text{NR}^1\text{R}^2$  and  $\text{R}^1$  and  $\text{R}^2$  are hydrogen may further be reacted with an acyl halide or an alkyl chloroformate in a reaction-inert solvent such as, for example dichloromethane, in the presence of a suitable base, such as, for example, pyridine, to form the corresponding amide, respectively, carbamate derivative.

Some of the compounds of formula (I') and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereochemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.


An alternative manner of separating the enantiomeric forms of the compounds of formula (I') and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

The above specified reaction procedures for the preparation of compounds of formula (I') or subgroups thereof, can also be applied for the preparation of compounds of formula (I).


Some of the intermediates and starting materials are known compounds and may be commercially available or may be prepared according to art-known procedures.

Intermediates of formula (II-A) wherein Q is  $\text{NR}^1\text{R}^2$ , said intermediates being represented by formula (II-A-1), can be prepared by reacting a pyrimidine derivative of formula (V) wherein  $\text{W}^2$  is a suitable leaving group such as, for example, a halogen, with  $\text{HNR}^1\text{R}^2$  (VI) in a reaction inert solvent such as, for example, 1,4-dioxane, 2-propanol or the like. Different regio-specific isomers may be formed and can be


separated from one another using suitable separation techniques such as, for example, chromatography.



Intermediates of formula (II-B) can be prepared analogously to the preparation of compounds of formula (I') starting from intermediates (II-A) and (III).




A particular subgroup of the intermediates of formula (II-B) is represented by formula




Particular intermediates of formula (II'-B) are those wherein  $\text{W}^1$  is a halogen, more in particular, a chloro atom.


Intermediates of formula (V) whereby Q is  $\text{NR}^1\text{R}^2$  and the L is  $\text{L}'\text{-CH}_2$  and is attached in the 2 position of the pyrimidine ring and  $\text{W}^2$  is chloro, said intermediates being represented by formula (V-a), can be prepared by reacting an imidamide of formula (VII) with a propanedioic acid ester of formula (VIII) in a solvent such as, for example, ethanol, and in the presence of, for instance, sodium, and subsequently reacting the thus formed intermediate of formula (IX) with a suitable reagent such as, for example, phosphoryl chloride.




Intermediates of formula (V) whereby Q is  $\text{NR}^1\text{R}^2$  and L is  $\text{L}'\text{-CH}_2$  and is attached in the 4 or 6 position of the pyrimidine ring and  $\text{W}^2$  is chloro, said intermediates being represented by formula (V-b), can be prepared by reacting an intermediate of formula (X) with urea or a functional derivative thereof, in a solvent such as, for example, ethanol, and in the presence of, for instance, sodium, and subsequently reacting the thus formed intermediate of formula (XI) with a suitable reagent such as, for example, phosphoryl chloride.



Intermediates of formula (V) wherein Q is  $\text{NR}^1\text{R}^2$  and L is  $\text{L}'\text{-CH}_2$  and is attached anywhere on the pyrimidine ring, said intermediates being represented by formula (V-c), can be prepared by reacting an intermediate of formula (XII-1), for Q is  $\text{NR}^1\text{R}^2$  and formula (XII-2) for Q is hydrogen, wherein  $\text{W}^2$  is a suitable leaving group such as, for example, a halogen, with an intermediate of formula (XIII) wherein  $\text{W}^3$  is a suitable leaving group such as, for example, a halogen, according to the procedure of a Grignard reaction.



Intermediates of formula (V) whereby Q is  $\text{NR}^1\text{R}^2$  and L is  $-\text{O}-\text{R}^6$  or  $-\text{NH}-\text{R}^6$  and is attached in the 4 or 6 position of the pyrimidine ring, said intermediates being represented by formula (V-d), can be prepared by reacting an intermediate of formula (XIV) with an intermediate of formula (XII) wherein  $\text{W}^2$  is a suitable leaving group such as, for example, a halogen, in a reaction-inert solvent such as, for example, tetrahydrofuran or 1,4-dioxane, and in the presence of a suitable base such as, for example, potassium hydroxide or diisopropyl ethaneamine, or sodium hydride.



The intermediates of formula (V-a) to (V-d) can analogously be prepared for the compounds of formula (I') wherein Q is hydrogen. To this effect, there is one leaving group  $W^2$  less on the pyrimidine ring of the respective starting material.

Compounds of formula (I') and some of the intermediates may have one or more stereogenic centers in their structure, present in a R or a S configuration.

The compounds of formula (I') as prepared in the hereinabove described processes may be synthesized as a mixture of stereoisomeric forms, in particular in the form of racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The compounds of the present invention and the intermediates of formula (II'-B) show antiretroviral properties, in particular against Human Immunodeficiency Virus (HIV), which is the aetiological agent of Acquired Immune Deficiency Syndrome (AIDS) in humans. The HIV virus preferentially infects human T-4 cells and destroys them or changes their normal function, particularly the coordination of the immune system. As a result, an infected patient has an everdecreasing number of T-4 cells, which moreover behave abnormally. Hence, the immunological defense system is unable to combat infections and neoplasms and the HIV infected subject usually dies by opportunistic infections such as pneumonia, or by cancers. Other conditions associated with HIV

infection include thrombocytopaenia, Kaposi's sarcoma and infection of the central nervous system characterized by progressive demyelination, resulting in dementia and symptoms such as, progressive dysarthria, ataxia and disorientation. HIV infection further has also been associated with peripheral neuropathy, progressive generalized lymphadenopathy (PGL) and AIDS-related complex (ARC).

The present compounds also show activity against HIV-1 strains that have acquired resistance to art-known non-nucleoside reverse transcriptase inhibitors. They also have little or no binding affinity to human  $\alpha$ -1 acid glycoprotein.

Due to their antiretroviral properties, particularly their anti-HIV properties, especially their anti-HIV-1-activity, the compounds of the present invention are useful in the treatment of individuals infected by HIV and for the prophylaxis of these individuals. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the enzyme reverse transcriptase. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic CNS diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

The compounds of the present invention or any subgroup thereof may therefore be used as medicines against above-mentioned conditions. Said use as a medicine or method of treatment comprises the systemic administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV and other pathogenic retroviruses, especially HIV-1.

The compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the

compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included.

Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions and the like, and segregated multiples thereof.

Those of skill in the treatment of HIV-infection could determine the effective daily amount from the test results presented here. In general it is contemplated that an effective daily amount would be from 0.01 mg/kg to 50 mg/kg body weight, more preferably from 0.1 mg/kg to 10 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

The exact dosage and frequency of administration depends on the particular compound

of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines and are not intended to limit the scope or use of the invention to any extent.

- Also, the combination of an antiretroviral compound and a compound of the present invention can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of the present invention, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in anti-HIV treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. Said other antiretroviral compounds may be known antiretroviral compounds such as nucleoside reverse transcriptase inhibitors, e.g. zidovudine (3'-azido-3'-deoxythymidine, AZT), didanosine (dideoxy inosine; ddI), zalcitabine (dideoxycytidine, ddC) or lamivudine (3'-thia-2'-3'-dideoxycytidine, 3TC) and the like; non-nucleoside reverse transcriptase inhibitors such as suramine, foscarnet-sodium (trisodium phosphono formate), nevirapine (11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido[3,2-b : 2',3'-e][1,4]diazepin-6-one), sustiva (efavirenz), tacrine (tetrahydroaminoacridine) and the like; compounds of the TIBO (tetrahydro-imidazo[4,5,1-jk][1,4]-benzodiazepine-2(1H)-one and thione)-type e.g. (S)-8-chloro-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-2-butenyl)imidazo-[4,5,1-jk][1,4]benzodiazepine-2(1H)-thione; compounds of the  $\alpha$ -APA ( $\alpha$ -anilino phenyl acetamide) type e.g.  $\alpha$ -[(2-nitro-phenyl)amino]-2,6-dichloro-benzene-acetamide and the like; TAT-inhibitors, e.g. RO-5-3335 and the like; protease inhibitors e.g. indinavir, ritanovir, saquinovir and the like; NMDA receptor inhibitors e.g. pentamidine;  $\alpha$ -glycosidase inhibitor e.g. castanospermine and the like; RNase H inhibitor e.g. dextran (dextran sulfate) and the like; or immunomodulating agents, e.g. levamisole, thymopentin and the like.

The following examples are intended to illustrate the present invention.

#### Experimental part

##### 35 A. Intermediate compounds

##### Example A1

a) A solution of 2,6-dichlorobenzylchloride (0.102 mol) in 1,1-diethylether (10 ml) was added dropwise to magnesium (0.102 mol) in 1,1-diethylether (60 ml). The

reaction was initiated by adding 2 drops of 1,2-dibromoethane. After most of magnesium disappeared, 2,4,6-trichloropyrimidine (0.051 mol) in 1,1-diethylether (30 ml) was added. The mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was purified by flash column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/hexane 1/2). The desired fractions were collected and the solvent was evaporated, yielding 3.3 g of (21%) 2,4-dichloro-6-[(2,6-dichloro-phenyl)methyl]pyrimidine (interm. 1; m.p.: 106-107 °C).

b) Intermediate (1) (0.0081 mol) in 2-propanol (100 ml) was heated until complete dissolution. The solution was then transferred into a pressure tube and NH<sub>3</sub> gas was bubbled into it for 20 minutes. Then the mixture was heated to 80 °C for 16 hours. The solvent was evaporated, yielding a residue of two compounds : 2-chloro-6-[(2,6-dichloro-phenyl)methyl]-4-pyrimidinamine (interm. 2) and 4-chloro-6-[(2,6-dichloro-phenyl)methyl]-2-pyrimidinamine (interm. 3).

#### Example A2

a) Urea (0.03 mol) was added to a mixture of (±)-ethyl 2,6-dichloro-phenyl-α-methyl-β-oxobutanoate (0.02 mol) in NaOC<sub>2</sub>H<sub>5</sub> in ethanol, (1M; 0.040 mol; 40 ml). The reaction mixture was stirred and refluxed overnight. The solvent was evaporated, water was added and the mixture was neutralized with 0.3 N HOAc. The precipitate was filtered off and was further triturated with ether and then H<sub>2</sub>O, then filtered off and dried, yielding 2.2 g (39%) of 6-[(2,6-dichloro-phenyl)methyl]-5-methyl-2,4(1*H*,3*H*)-pyrimidinedione (interm. 4).

b) A mixture of intermediate (4) (0.0095 mol) in phosphoryl chloride (50 ml) was stirred and refluxed overnight. Excess of phosphoryl chloride was then evaporated. Ice-water was added to the residue. A white precipitate was formed, filtered off and dried. The residue was purified by flash column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>). The desired fractions were collected and the solvent was evaporated, yielding 2.06 g (67%) of 2,4-dichloro-6-[(2,6-dichloro-phenyl)methyl]-5-methyl-pyrimidine (interm. 5).

c) 4-chloro-6-[(2,6-dichloro-phenyl)methyl]-5-methyl-2-pyrimidinamine (interm. 6) and 2-chloro-6-[(2,6-dichloro-phenyl)methyl]-5-methyl-4-pyrimidinamine (interm. 7) were prepared from intermediate 5 following the procedures as described in example A1b.

#### Example A3

a) To the stirred solution of 2,6-dichlorobenzeneethanimidamide HCl (1:1), (0.0042 mol) in ethanol (20 ml), a solution of sodium (0.013 mol) in ethanol (10 ml) was added dropwise first and then propanedioic acid, diethyl ester (0.0109 mol) was added. The

reaction mixture was stirred and refluxed for 4 hours and then stirred at room temperature overnight. After adding another equivalent of propanedioic acid, diethyl ester (stirring and refluxing it overnight), the solvent was evaporated and the residue was dissolved in water and acidified with 1 N HCl. The solid was filtered off, washed with water and dried, yielding 0.87 g (76.4%) of 2-[(2,6-dichloro-phenyl)methyl]-4,6-pyrimidinediol (interm. 8).

b) 6-chloro-2-[(2,6-dichloro-phenyl)methyl]-4-pyrimidinamine (interm. 9) was prepared starting from intermediate 8 according to the procedures described in example A.1.b, A2.b & A2.c.

#### Example A4

4-Amino-1-butanol (1.57 ml) was added to a solution of intermediate (1) (0.008 mol) in 1,4-dioxane (20 ml) under Argon. The reaction mixture was stirred for 2 hours at room temperature. The solvent was evaporated. The residue was purified by flash column chromatography over silica gel (eluent gradient: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH: from 100/0 to 98/2). The pure fractions were collected and the solvent was evaporated, yielding 2.05 g of a mixture of 4-[[2-chloro-6-[(2,6-dichloro-phenyl)methyl]-4-pyrimidinyl]-amino]-1-butanol (interm. 10) and 4-[[4-chloro-6-[(2,6-dichloro-phenyl)methyl]-2-pyrimidinyl]amino]-1-butanol (interm. 11).

#### Example A5

a) Potassium hydroxide/ethanol (10%; 0.035 mol) was added to a solution of 2,6-dichlorophenol (0.035 mol) in tetrahydrofuran (100 ml). The mixture was stirred and 2,4,6-trichloropyrimidine (0.044 mol) was added. The mixture was stirred overnight at 60 °C. The reaction was quenched with NaOH 1N solution. The aqueous layers were extracted with EtOAc several times and then the organic layers were combined and washed with NaOH 3N and saturated NaCl, dried and concentrated. The residue was recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/hexane. The precipitate was filtered off and dried, yielding 5.98 g 2,4-dichloro-6-(2,6-dichlorophenoxy)pyrimidine (55%) (interm. 12).

b) Reaction under Argon atmosphere. 2,4,6-trimethylaniline (0.0678 mol) was added to 2,4-dichloropyrimidine (0.0664 mol) in 1,4-dioxane (100 ml). *N,N*-di(1-methylethyl)-ethaneamine (0.0830 mol) was added. The reaction mixture was stirred and refluxed for 4 days and the solvent was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, washed with a saturated NaHCO<sub>3</sub> solution, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and the solvent was evaporated to give 17.1 g solid residue. This solid was dissolved in CH<sub>2</sub>Cl<sub>2</sub>:hexane (1:1; 150 ml), and the resulting solution was concentrated to 100 ml, then filtered. The residue was purified by column chromatography on KP-Sil (eluent: CH<sub>2</sub>Cl<sub>2</sub>). The desired fractions were collected and the solvent was evaporated. The less polar fraction

was stirred in CH<sub>2</sub>Cl<sub>2</sub> for 3 hours and filtered, yielding 0.44 g 4-chloro-*N*-(2,4,6-trimethylphenyl)-2-pyrimidinamine (intermediate 48). A second fraction was recrystallized from acetonitrile, filtered off and dried, yielding 2-chloro-*N*-(2,4,6-trimethyl-phenyl)-4-pyrimidinamine (intermediate 49).

5 Example A6

Pyridine (1 ml) was added to a mixture of 4-[[4-amino-6-[(2,6-dichloro-phenyl)-methyl]-2-pyrimidinyl]amino]benzonitrile (0.00135 mol) in CH<sub>2</sub>Cl<sub>2</sub> (19 ml). A solution of chloroethanoyl chloride (0.001375 mol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 ml) was added dropwise on an ice bath. The mixture was stirred at room temperature for 2 hours.

- 10 More chloroethanoyl chloride (0.00625 mol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 ml) was added. The mixture stood in the refrigerator overnight. The solvent was evaporated. The residue was treated with a saturated Na<sub>2</sub>CO<sub>3</sub> solution and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The separated organic layer was dried, filtered and concentrated. The residue was purified by flash column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/  
15 NH<sub>4</sub>OH 99/1/0.1). The desired fractions were collected and the solvent was evaporated, yielding 0.22 g (36.5%) of 2-chloro-*N*-[6-[(2,6-dichloro-phenyl)methyl]-2-[(4-cyano-phenyl)amino]-4-pyrimidinyl]acetamide (interm. 13).

Example A7


- A mixture of 4-[(4-chloro-2-pyrimidinyl)amino]benzonitrile (0.005 mol) and nityl  
20 tetrafluoroborate (0.0025 mol) in acetonitrile (5 ml) was stirred at room temperature for 4 h. The material was quenched with saturated bicarbonate (50 ml) on cracked ice. The mixture was allowed to reach room temperature, and the yellow solid was filtered off. The solid was adsorbed onto silica and purified by column chromatography (eluent: 30%, 50%, 60%, 70% CH<sub>2</sub>Cl<sub>2</sub> in hexanes). The solvent of the desired fraction was  
25 evaporated and the residue was dried, yielding 0.89 g (64%) of 3-nitro-4-[(4-chloro-2-pyrimidinyl)amino]benzonitrile.(interm. 51)

Example A8

- A mixture of 2,6-dichloro-*N*-(2,4,6-trimethylphenyl)-4-pyrimidinamine (0.00376 mol) in a 2.0 M solution of NH<sub>3</sub> in 2-propanol (25 ml) and a 0.5 M solution of NH<sub>3</sub> in  
30 dioxane (25 ml) was heated in a pressure sample at 110-115 °C for 24 hours. The solvent was evaporated, and the residue was chromatographed on Biotage (eluent: 1:1 CH<sub>2</sub>Cl<sub>2</sub>:hexane). The desired fractions were collected and the solvent was evaporated, yielding a mixture of 0.523 g of 2-chloro-*N*4-(2,4,6-trimethylphenyl)-4,6-pyrimidine-  
35 pyrimidinediamine (interm. 53) and 0.101 g of 6-chloro-*N*4-(2,4,6-trimethylphenyl)-2,4-pyrimidinediamine. (interm. 50)

Tables 1 and 2 list intermediates which were prepared analogous to one of the above

examples.

Table 1a





Int. No.	Ex. No.	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	X	R <sup>5</sup>	R	physical data melting point
6	A2c	Cl	H	Cl	CH <sub>2</sub>	CH <sub>3</sub>	-NH <sub>2</sub>	-
15	A1b	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>3</sub>	-
16	A1b	Cl	H	Cl	O	H	-NH-CH <sub>3</sub>	152-155°C
17	A1b	Cl	H	Cl	O	H	-NH <sub>2</sub>	-
19	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>3</sub> -OH	-
20	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -OH	111-113°C
21	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>2</sub> -CH(OH)-C <sub>6</sub> H <sub>5</sub>	133-134°C
22	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>3</sub> -N <sub>1</sub> 	-
23	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -O-(CH <sub>2</sub> ) <sub>2</sub> OH	99-107°C
24	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(4-OCH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )	138-140°C
25	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(3-OCH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )	132-135°C
26	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>2</sub> -CH(OH)-CH <sub>2</sub> OH	116-118°C
27	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub>	137-139°C
28	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(2-thienyl)	113-114°C
29	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(2-pyridyl)	113.5-114°C
31	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> CN	151-153°C
48	A5b	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	NH	H	-H	142-143°C
50	A8	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	NH	H	-NH <sub>2</sub>	

Table 1b

Int. No.	Ex. No.	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	X	R <sup>5</sup>	R	physical data melting point
14	A2b	H	CN	H	NH	H	H	211-212°C
18	A5b	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	NH	CH <sub>3</sub>	H	
30	A2b	H	CN	H	NH	CH <sub>3</sub>	H	
51	A7	NO <sub>2</sub>	CN	H	NH	H	H	
								142-144°C

Table 2



Int. No.	Ex. No.	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	X	R <sup>5</sup>	R	physical data
7	A2c	Cl	H	Cl	CH <sub>2</sub>	CH <sub>3</sub>	-NH <sub>2</sub>	134-135°C 130-133°C 98-99°C 104-109°C 149-150°C 137-139°C - 170-173°C - - 171-172°C >60°C 137-141°C 55-60°C  182-183°C 75-83°C 159-161°C
32	A1b	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>3</sub>	
33	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(1-pyrrolidiny)	
34	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(2-pyridyl)	
35	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(2-thienyl)	
36	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(3-OCH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )	
37	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(4-OCH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )	
38	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> CN	
39	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> -O-(CH <sub>2</sub> ) <sub>2</sub> OH	
40	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>2</sub> OH	
41	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>3</sub> -O-CH(CH <sub>3</sub> ) <sub>2</sub>	
42	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>3</sub> -OH	
43	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub>	
45	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>2</sub> -CH(OH)-CH <sub>2</sub> OH	
46	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-O-CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub>	
47	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-(CH <sub>2</sub> ) <sub>3</sub> -N 	
49	A5b	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	NH	H	H	
52	A4	Cl	H	Cl	CH <sub>2</sub>	H	-NH-CH <sub>2</sub> -CH(OH)-C <sub>6</sub> H <sub>5</sub>	
53	A1b	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	NH	H	-NH <sub>2</sub>	
54	A5b	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	NH	CH <sub>3</sub>	H	
55	A5a	Cl	Cl	Cl	-O-	H	H	

## B. Compounds of formula (I')

### Example B1

A mixture of intermediate (42) and intermediate (19) (0.004 mol) and 4-amino-benzonitrile (0.0084 mol) were combined in a sealed tube and heated for 16 hours at 160°C under Argon. The reaction mixture was allowed to cool to room temperature and dissolved in CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 90/10 (20 ml) and 5 g of silica gel was added. After evaporating the solvent, the residue was purified by flash column chromatography over silica gel (eluent gradient: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH: from 100/0 to 97/3). The desired fraction was collected and the solvent was evaporated, yielding 0.31 g (18.1%) of 4-[[4-[(2,6-dichloro-phenyl)methyl]-6-[(3-hydroxypropyl)amino]-2-pyrimidinyl]amino]benzonitrile (compound 3).

### Example B2

Intermediates (47) and (22) (0.00399 mol) and 4-aminobenzonitrile (0.0012 mol) in 1-methyl-2-pyrrolidinone (3 ml) was stirred for 16 hours at 130 °C under Argon. Then, the reaction mixture was cooled to room temperature and quenched with H<sub>2</sub>O (200 ml). A precipitate formed, which was stirred for 16 hours, and separated by filtration over Celite. The residue was dissolved in CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> (10%, 200 ml), dried over K<sub>2</sub>CO<sub>3</sub>, filtered, and evaporated. This resulting material was further purified by flash column chromatography over silica gel (gradient eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH from 100/0 to 95/5). The desired fraction was collected and the solvent was evaporated, yielding 0.43 g (21.7%) of 4-[[6-[(2,6-dichloro-phenyl)methyl]-2-[[3-(2-oxo-1-pyrrolidinyl)propyl]-amino]-4-pyrimidinyl]amino]benzonitrile (comp. 39; 104-114°C).

### Example B3

HCl/diethyl ether (1N; 2.77 ml) was stirred into a solution of intermediate (33) (0.00277 mol) in 1-methyl-2-pyrrolidinone (4 ml) under N<sub>2</sub> atmosphere. The reaction mixture was heated for 5 minutes. Next, 4-aminobenzonitrile (0.0061 mol) was added and the reaction was heated at 100 °C for 16 hours. Then, the reaction mixture was cooled to room temperature and diluted with ethylacetate (10 ml). The organic layer was washed with NaOH (1 N; 2 x 100 ml), H<sub>2</sub>O (2 x 100 ml), brine (50 ml), respectively, dried, filtered and the filtrate was evaporated. The crude material was purified by flash chromatography (eluent: 2.5-7.5% of CH<sub>3</sub>OH containing 10% NH<sub>4</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>). The desired fractions were collected and the solvent was evaporated. The residue was dried, yielding 0.160 g (12.0%) of 4-[[4-[(2,6-dichloro-phenyl)methyl]-6-[[2-(1-pyrrolidinyl)ethyl]amino]-2-pyrimidinyl]amino]benzonitrile (comp. 13; mp. 80-85°C).

Example B4

A slurry of intermediate (14) (0.005 mol) in  $\text{CH}_2\text{Cl}_2$  (150 ml) was stirred rapidly and cooled to 0 °C under nitrogen.  $\text{BBr}_3$  (0.015 mol) was introduced by syringe. The reaction mixture was stirred rapidly for two hours. The reaction mixture was recooled to 0 °C and quenched with NaOH (aq. 1 N, 25 ml). The biphasic partial quench mixture gives a precipitate which was filtered off and dried, yielding 2.5 g (91%) of 4-[[4-[(2,6-dichloro-phenyl)methyl]-6-(hydroxyamino)-2-pyrimidinyl]amino]-benzonitrile dihydrobromide.pentahydrate (comp. 15; mp. 240-244°C).

Example B5

1,1-Dimethoxy-*N,N*-dimethylmethanamine (0.152 mol) was added to 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]benzonitrile (0.0008 mol). The mixture was stirred at room temperature for 2 days and then concentrated. The crude product was purified by flash chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1). The desired fraction was collected and the solvent was evaporated. The resulting residue was triturated with hexane, yielding 0.15 g (42%) of *N*'-[2-[(4-cyano-phenyl)amino]-6-[(2,6-dichlorophenyl)methyl]-4-pyrimidinyl]-*N,N*-dimethyl-methanimidamide (comp. 26; mp. 175-180°C).

Example B6

Piperidine (0.12 ml) was added to a mixture of intermediate (13) (0.00047 mol) in tetrahydrofuran (20 ml). The mixture was stirred at room temperature for 4 hours. More piperidine (0.14 ml) was added. The mixture was stirred for another 2 hours. The solvent was evaporated. The residue was purified by flash column chromatography over silica gel ( $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  99/1/0.1). The desired fractions were collected and the solvent was evaporated, yielding 0.05 g (21.5%) of *N*-[6-[(2,6-dichloro-phenyl)methyl]-2-[(4-cyano-phenyl)amino]-4-pyrimidinyl]-1-piperidine-acetamide (comp. 25; mp. 175-180°C).

Example B7

Pyridine (0.014 mol) was added to a mixture of 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]benzonitrile (0.0013 mol) in  $\text{CH}_2\text{Cl}_2$ . A solution of octanoyl chloride (1.5 equiv) in  $\text{CH}_2\text{Cl}_2$  (0.5 ml) was added dropwise. The mixture was stirred at room temperature for 2 hours. More octanoyl chloride (3.5 equiv) in  $\text{CH}_2\text{Cl}_2$  was added dropwise. The mixture was stirred. The solvent was then evaporated. The residue was treated with a saturated aqueous  $\text{NaHCO}_3$  solution and the mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The separated organic layer was dried, filtered and the solvent was evaporated to give the crude product. The residue was recrystallized from  $\text{CHCl}_3$  and hexane, yielding 0.443 g (68.6%) of *N*-[6-[(2,6-dichloro-phenyl)methyl]-2-[(4-

cyano-phenyl)amino]-4-pyrimidinyl]octanamide (comp. 17; mp. 135-137°C).

#### Example B8

5 a) A mixture of intermediate 49 (0.082 mol) and 5.4 N HCl in 2-propanol (0.086 mol) in water (300 ml) was stirred and warmed to 40-45 °C over 30 minutes. 4-Amino-  
benzonitrile (0.242 mol) was added at 40-45 °C. The reaction mixture was stirred and  
10 refluxed for 4.5 hours, then cooled to room temperature. The mixture was alkalized by  
portionwise addition of NaHCO<sub>3</sub>. This mixture was extracted with ethylacetate. The  
organic layer was separated, washed with brine, dried, filtered and the solvent was  
evaporated. This fraction was stirred in ethanol p.a. (100 ml), filtered off, washed with  
10 ethanol (50 ml), then dried, yielding 23.1 g (86%) 4-[[4-[(2,4,6-trimethylphenyl)-  
amino]-2-pyrimidinyl]amino]benzonitrile (compound 52).

b) A mixture of 4-[(4-chloro-2-pyrimidinyl)amino]benzonitrile (0.021 mol) and HCl in  
2-propanol (0.0095 mol) in water (30 ml) was stirred for one hour at 45 °C. 4-amino-  
3,5-dimethyl-benzonitrile (0.025 mol) was added and the reaction mixture was stirred  
15 and refluxed overnight. The mixture was cooled to room temperature, then neutralized  
with NaHCO<sub>3</sub>. This mixture was extracted with ethylacetate. The separated organic  
layer was washed with brine, dried, filtered and the solvent evaporated. The residue was  
crystallized from CH<sub>3</sub>CN, filtered off and dried. The residue was stirred in boiling  
CH<sub>2</sub>Cl<sub>2</sub> (20 ml), then filtered off and dried. The residue was crystallized from methyl  
20 isobutyl keton, filtered off and dried, yielding 0.3 g of 4-[[2-[(cyanophenyl)amino]-4-  
pyrimidinyl]amino]-3,5-dimethylbenzonitrile (compound 69).

#### Example B9

a) 4-[(4-chloro-2-pyrimidinyl)amino]benzonitrile (0.003 mol), 2,6-dibromo-4-methyl-  
benzenamine (0.006 mol) and 1 M HCl in diethyl ether (4.5 ml) in 1,4-dioxane (10 ml)  
25 were combined in a tube and heated under Ar until all diethyl ether had evaporated. The  
tube was sealed and heated at 170 °C for 2.5 days. Silica gel was added, and the solvent  
was evaporated. The residue was purified by flash column chromatography over silica  
gel (eluent gradient: CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH:NH<sub>4</sub>OH 100:0:0 to 99:0.9:0.1). The desired  
fractions were collected and the solvent was evaporated. The residue was recrystallized  
30 from acetonitrile, filtered off and dried, yielding 0.22 g (15.9%) of 4-[[4-[(2,6-dibromo-  
4-methylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile (compound 61).

b) 4-[[4-[(4-chloro-5-methyl-2-pyrimidinyl]amino]benzonitrile (0.01541 mol),  
4-amino-3,5-dimethyl-benzonitrile (0.00219 mol), 1-methyl-2-pyrrolidinone (4 ml),  
1,4-dioxane (15 ml) and diisopropylethylamine (0.0154 mol) were combined in a flask  
35 under a stream of argon and heated at 160-230 °C for 16 hours. CH<sub>2</sub>Cl<sub>2</sub> and 1N NaOH  
were added, and the mixture was stirred 1 hour and filtered to give a brown solid (\*).  
The CH<sub>2</sub>Cl<sub>2</sub> filtrate was separated and was evaporated and purified by flash column

chromatography (eluent: 2% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>). The desired fractions were combined, evaporated and the residue was stirred in CH<sub>2</sub>Cl<sub>2</sub>. The solid precipitate was filtered off, combined with the brown solid (\*) and recrystallized from CH<sub>3</sub>CN. The precipitate was filtered off and dried, yielding 1.57 g (29%) of 4-[[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile (compound 89).

c) 2-[(4-cyanophenyl)amino]-4-pyrimidinyl trifluoromethanesulfonate (0.0022 mol) and 2,6-dichloro-4-(trifluoromethyl)-benzenamine (0.0044 mol) were combined in 1,4-dioxane (2.5 ml) and heated in a sealed tube under Ar at 170 °C for 40 hours. The reaction mixture was allowed to cool to room temperature. Silica gel was added, and the solvent was evaporated. The residue was purified by flash column chromatography over silica gel (eluent gradient: CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH:NH<sub>4</sub>OH 100:0:0 to 97:2.7:0.3). The desired fractions were collected and the solvent was evaporated. The residue was recrystallized from CH<sub>3</sub>CN, filtered off and dried, yielding 0.086 g (9.2%) of 4-[[4-[[2,6-dichloro-4-(trifluoromethyl)-phenyl]amino]-2-pyrimidinyl]amino]benzonitrile (compound 66).

#### Example B10

To a suspension of NaH (0.006 mol) in 1,4-dioxane (30 ml), 2,4,6-trimethyl-phenol (0.006 mol) was added. The mixture was stirred for 15 minutes at room temperature, and a clear solution formed. 4-[(4-chloro-2-pyrimidinyl)amino]benzonitrile (0.004 mol) was added, and the reaction mixture was heated to reflux under Argon for 15 hours. The reaction mixture was allowed to cool to room temperature, 0.5 ml of water was added, followed by 4 g of silica gel, and the solvent was evaporated. The residue was purified by flash column chromatography over silica gel (eluent gradient: CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH 100:0:0 to 97:3). The pure fractions were collected and the solvent was evaporated, yielding 1.18 g (89.4%) of 4-[[4-(2,4,6-trimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile (compound 58).

#### Example B11

Compound (52) (0.0015 mol) was stirred in boiling ethanol (8 ml). 6 M HCl in 2-propanol (0.0015 mol) was added and the salt was allowed to crystallize out overnight at room temperature. The precipitate was filtered off, washed with 2-propanol and dried, yielding 0.47 g (86%) of 4-[[4-[(2,4,6-trimethyl-phenyl)amino]-2-pyrimidinyl]amino]benzonitrile hydrochloride (1:1) (compound 53).

#### Example B12


A mixture of compound (52) (0.00303 mol) and NaBO<sub>3</sub>·4H<sub>2</sub>O (0.00911 mol) in CH<sub>3</sub>OH (30 ml) and H<sub>2</sub>O (10 ml) was stirred and refluxed for 4 days. The reaction mixture was

cooled. The precipitate was filtered off and the precipitate (\*) was purified by flash column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  gradient from 100/0 to 95/5). The desired fractions were collected and the solvent was evaporated, yielding 0.586 g (56%) of 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-

5 pyrimidinyl]amino]benzamide (compound 100). The filtrate (\*) was purified by reversed-phase HPLC (eluent gradient: ((0.5% ammoniumacetate in  $\text{H}_2\text{O}$ )/ $\text{CH}_3\text{CN}$  90/10)/ $\text{CH}_3\text{OH}/\text{CH}_3\text{CN}$  (0 minutes) 75/25/0, (44 minutes) 0/50/50, (57 minutes) 0/0/100, (61.1-70 minutes) 75/25/0). Three desired fraction groups were collected and their solvent was evaporated, yielding 0.18 g of 4-[[4-[(2,4,6-trimethylphenyl)amino]-  
10 2-pyrimidinyl]amino]benzamide, *N*3-oxide (compound 106) and 0.030 g of 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzamide, *N*1-oxide (compound 107).

Tables 3, 4, 5 and 6 list the compounds of formula (I) that were prepared according to one of the above examples.


15 Table 3



Co. No.	Ex. No.	$\text{NR}^1\text{R}^2$	physical data (melting point in $^{\circ}\text{C}$ )
1	B2	$-\text{NH}-(\text{CH}_2)_4-\text{OH}$	161-163 $^{\circ}\text{C}$
2	B2	$-\text{NH}-(\text{CH}_2)_2-\text{OH}$	207-210 $^{\circ}\text{C}$
3	B2	$-\text{NH}-(\text{CH}_2)_3-\text{OH}$	152-154 $^{\circ}\text{C}$
4	B2	$-\text{NH}-\text{CH}_2-\text{CHOH}-\text{C}_6\text{H}_5$	158-165 $^{\circ}\text{C}$
5	B2	$-\text{NH}-(\text{CH}_2)_3-\text{N} \begin{array}{c} \diagup \diagdown \\ \text{ } \end{array} \begin{array}{c} \diagup \diagdown \\ \text{ } \end{array} \text{O}$	48-56 $^{\circ}\text{C}$
6	B2	$-\text{NH}-(\text{CH}_2)_2-\text{O}-(\text{CH}_2)_2-\text{OH}$	162-175 $^{\circ}\text{C}$ ; HCl (1:1)
7	B2	$-\text{NH}-(\text{CH}_2)_3-\text{O}-\text{CH}(\text{CH}_3)_2$	181-182 $^{\circ}\text{C}$ ; HCl (1:1)
8	B2	$-\text{NH}-(\text{CH}_2)_2-(3-\text{OCH}_3-\text{C}_6\text{H}_4)$	72-80 $^{\circ}\text{C}$
9	B2	$-\text{NH}-\text{CH}_2-\text{CHOH}-\text{CH}_2\text{OH}$	189-192 $^{\circ}\text{C}$
10	B2	$-\text{NH}-(\text{CH}_2)_2-(4-\text{OCH}_3-\text{C}_6\text{H}_4)$	72-80 $^{\circ}\text{C}$

Co. No.	Ex. No.	NR <sup>1</sup> R <sup>2</sup>	physical data (melting point in °C)
11	B2	-NH-O-CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub>	-
12	B2	-NH-CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub>	-
13	B3	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(1-pyrrolidiny)	80-85°C
14	B2	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(2-thienyl)	-
15	B4	-NH-OH	240-244°C
16	B2	-NH-(CH <sub>2</sub> ) <sub>2</sub> -(2-pyridyl)	75-80°C
17	B7	-NH-CO-C <sub>7</sub> H <sub>15</sub>	135-137°C
18	B7	-NH-CO-C <sub>11</sub> H <sub>23</sub>	130-135°C
19	B2	-NH-(CH <sub>2</sub> ) <sub>2</sub> -CN	255°C; HCl (1:1)
20	B7	-NH-CO-O-C <sub>2</sub> H <sub>5</sub>	>200°C
21	B7	-NH-CO-CH <sub>3</sub>	128-130°C
22	B7	-NH-CO-C <sub>3</sub> H <sub>7</sub>	>200°C
23	B1	-NH <sub>2</sub>	94-97°C
24	B1	-NH-CH <sub>3</sub>	178-180°C
25	B6	-NH-CO-CH <sub>2</sub> -(1-piperidiny)	175-180°C
26	B5	-N=CH-N(CH <sub>3</sub> ) <sub>2</sub>	175-180°C

Table 4



Co. No.	Ex. No.	R'	R''	R <sup>5</sup>	physical data (melting point)
27	B1	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	-
28	B1	H	4-Br-C <sub>6</sub> H <sub>4</sub>	H	-
29	B1	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	-
30	B1	H	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	-
31	B1	H	(3-Br-6-pyridyl)	H	-
32	B1	(3-Br-6-pyridyl)	H	H	-
33	B1	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	77-80°C
34	B1	H	4-F-C <sub>6</sub> H <sub>4</sub>	H	>200°C
35	B1	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	76-79°C
36	B1	H	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	183-186°C




Co. No.	Ex. No.	R'	R''	R <sup>5</sup>	physical data (melting point)
37	B1	C <sub>6</sub> H <sub>5</sub>	H	H	85-90°C
38	B1	H	C <sub>6</sub> H <sub>5</sub>	H	182-187°C
39	B2		4-CN-C <sub>6</sub> H <sub>4</sub>	H	104-114°C
40	B2	(CH <sub>2</sub> ) <sub>2</sub> -OH	4-CN-C <sub>6</sub> H <sub>4</sub>	H	247-250°C; HCl (1:1)
41	B1	CH <sub>3</sub>	4-CN-C <sub>6</sub> H <sub>4</sub>	H	>200°C
42	B1	(CH <sub>2</sub> ) <sub>3</sub> -OH	4-CN-C <sub>6</sub> H <sub>4</sub>	H	91-105°C
43	B2	(CH <sub>2</sub> ) <sub>4</sub> -OH	4-CN-C <sub>6</sub> H <sub>4</sub>	H	161-163°C
45	B1	H	4-CN-C <sub>6</sub> H <sub>4</sub>	H	>200°C
46	B1	H	4-CN-C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	>200°C
47	B1	4-CN-C <sub>6</sub> H <sub>4</sub>	H	CH <sub>3</sub>	>200°C
48	B1	H	4-Br-C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	>200°C
49	B1	4-Br-C <sub>6</sub> H <sub>4</sub>	H	CH <sub>3</sub>	168-170°C

Table 5



Co. No.	Ex. No.	R'	R''	R'''	R <sup>5</sup>	physical data
50	B1	NH <sub>2</sub>	4-CN-C <sub>6</sub> H <sub>4</sub>	O-(2,6-diCl-C <sub>6</sub> H <sub>3</sub> )	H	>200°C
51	B1	CH <sub>2</sub> -(2,6-diCl-C <sub>6</sub> H <sub>3</sub> )	H	-NH-(4-CN-C <sub>6</sub> H <sub>4</sub> )	H	>200
90	B9a	NH-(2-NO <sub>2</sub> -4-CN-C <sub>6</sub> H <sub>3</sub> )	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	165-168°C
91		NH-(3-OH-4-CN-C <sub>6</sub> H <sub>3</sub> )	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
92	B12	NH-(2,6-diCl-C <sub>6</sub> H <sub>3</sub> )	2,6-diCl-C <sub>6</sub> H <sub>3</sub>	H	H	164-166°C
93	B9a	NH-(2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub> )	4-CN-C <sub>6</sub> H <sub>4</sub>	H	H	267-268°C
94	B1	NH-(4-CN-C <sub>6</sub> H <sub>4</sub> )	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	NH <sub>2</sub>	H	263-264°C
95	B1	NH <sub>2</sub>	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	-NH-(4-CN-C <sub>6</sub> H <sub>4</sub> )	H	233-234°C
96	B8a	NH-(4-Cl-C <sub>6</sub> H <sub>4</sub> )	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
97	B8a	NH-(2,4-diF-C <sub>6</sub> H <sub>3</sub> )	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
98	B8a		2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
99	B9a	NH-(2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub> )	4-CN-C <sub>6</sub> H <sub>4</sub>	H	CH <sub>3</sub>	200-201°C






Co. No.	Ex. No.	R'	R''	R'''	R <sup>5</sup>	physical data
100	B11		2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
101	B8a		2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
102	B8a		2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	
103	B1		H	-CH <sub>2</sub> -(2,6-diCl-C <sub>6</sub> H <sub>3</sub> )	CH <sub>3</sub>	>200°C
104		NH-(4-CN-C <sub>6</sub> H <sub>4</sub> )	C <sub>6</sub> H <sub>5</sub> -CH <sub>2</sub> -	H	H	
105		NH-(2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub> )	2,4,6-triCH <sub>3</sub> -C <sub>6</sub> H <sub>2</sub>	H	H	

Table 6



Co. No.	Ex. No.	X	R <sup>5</sup>	R <sup>6a</sup>	R <sup>6b</sup>	R <sup>6c</sup>	R <sup>6d</sup>	physical data (salt form; melting point)
52	B8a	NH	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	217-218°C
53	B11	NH	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	HCl (1:1)
54	B11	NH	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	HBr (1:1)
55	B11	NH	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	L-tartrate
56	B9a	NH	H	CH <sub>3</sub>	H	Br	CH <sub>3</sub>	HCl (1:1); 214-217°C
57	B9a	NH	H	CH <sub>3</sub>	H	H	CH <sub>3</sub>	HCl (1:1); > 270°C
58	B10	O	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	220-222°C
59	B10	S	H	Cl	H	H	Cl	225-226°C
60	B3	O	H	Cl	H	Cl	Cl	279-280°C
61	B9a	NH	H	Br	H	CH <sub>3</sub>	Br	230-233°C
62	B9a	NH	H	Br	H	CH(CH <sub>3</sub> ) <sub>2</sub>	Br	198-200°C

Co. No.	Ex. No.	X	R <sup>5</sup>	R <sup>6a</sup>	R <sup>6b</sup>	R <sup>6c</sup>	R <sup>6d</sup>	physical data (salt form; melting point)
63	B3	NH	CH <sub>3</sub>	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	236-237°C
64	B10	O	H	Cl	H	Cl	CH <sub>3</sub>	266-267°C
65	B9a	NH	H	Cl	H	H	Cl	253-255°C
66	B9c	NH	H	Cl	H	CF <sub>3</sub>	Cl	239-240°C
67	B9c	NH	H	Br	H	F	Cl	244-245°C
68	B9a	NH	H	Cl	H	Cl	CH <sub>3</sub>	217°C
69	B8b or B9a	NH	H	CH <sub>3</sub>	H	CN	CH <sub>3</sub>	225-230°C
70	B9c	NH	H	Br	H	Br	F	210-214°C
71	B9c	N(CH <sub>3</sub> )	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	218-219°C
72	B9c	NH	H	Cl	H	Cl	Cl	trifluoroacetate (1:1); 225-230°C
73	B10	S	H	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	204.5-208°C
74	B10	O	H	Br	H	Cl	CH <sub>3</sub>	246-249°C
75	B9c	NH	H	CH <sub>3</sub>	H	Cl	CH <sub>3</sub>	206-207°C
76	B9a	NH	H	Cl	H	CN	Cl	>180°C
77	B9c	NH	H	Cl	H	OCF <sub>3</sub>	Cl	185-190°C
78	B9c	NH	H	Br	Cl	Br	Cl	>265°C
79	B9c	NH	H	Br	H	C <sub>3</sub> H <sub>7</sub>	Br	215-218°C
80	B9a	NH	H	CH <sub>3</sub>	H	C(CH <sub>3</sub> ) <sub>3</sub>	CH <sub>3</sub>	203-205°C
81	B10	O	H	CH <sub>3</sub>	H	CN	CH <sub>3</sub>	279-280°C
82	B9c	NH	CH <sub>3</sub>	CH <sub>3</sub>	H	Cl	CH <sub>3</sub>	235-237°C
83	B9b	NH	CH <sub>3</sub>	CH <sub>3</sub>	H	CN	CH <sub>3</sub>	H <sub>2</sub> O (1:1) trifluoroacetate (1:1); 274-275°C
84	B9c	NH	CH <sub>3</sub>	CH <sub>3</sub>	H	C(CH <sub>3</sub> ) <sub>3</sub>	CH <sub>3</sub>	231-232°C
85	B9c	NH	CH <sub>3</sub>	CH <sub>3</sub>	H	Br	CH <sub>3</sub>	218-219°C
86	B9c	S	CH <sub>3</sub>	CH <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	229-230°C
87	B9a	NH	CH <sub>3</sub>	Br	H	C <sub>3</sub> H <sub>7</sub>	Br	197-198°C
88	B9a	NH	CH <sub>3</sub>	Br	H	CH(CH <sub>3</sub> ) <sub>2</sub>	Br	157-158°C
89	B9b	NH	CH <sub>3</sub>	CH <sub>3</sub>	H	CN	CH <sub>3</sub>	>300°C

### C. Pharmacological example

#### Example C.1

A rapid, sensitive and automated assay procedure was used for the *in vitro* evaluation of anti-HIV agents. An HIV-1 transformed T4-cell line, MT-4, which was previously shown (Koyanagi et al., *Int. J. Cancer*, **36**, 445-451, 1985) to be highly susceptible to

and permissive for HIV infection, served as the target cell line. Inhibition of the HIV-induced cytopathic effect was used as the end point. The viability of both HIV- and mock-infected cells was assessed spectrophotometrically via the *in situ* reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT). The 50% cytotoxic concentration (CC<sub>50</sub> in  $\mu\text{M}$ ) was defined as the concentration of compound that reduced the absorbance of the mock-infected control sample by 50%. The percent protection achieved by the compound in HIV-infected cells was calculated by the following formula :

$$\frac{(\text{OD}_T)_{\text{HIV}} - (\text{OD}_C)_{\text{HIV}}}{(\text{OD}_C)_{\text{MOCK}} - (\text{OD}_C)_{\text{HIV}}} \quad \text{expressed in \%},$$

whereby (OD<sub>T</sub>)<sub>HIV</sub> is the optical density measured with a given concentration of the test compound in HIV-infected cells; (OD<sub>C</sub>)<sub>HIV</sub> is the optical density measured for the control untreated HIV-infected cells; (OD<sub>C</sub>)<sub>MOCK</sub> is the optical density measured for the control untreated mock-infected cells; all optical density values were determined at 540 nm. The dose achieving 50% protection according to the above formula was defined as the 50% inhibitory concentration (IC<sub>50</sub> in  $\mu\text{M}$ ). The ratio of CC<sub>50</sub> to IC<sub>50</sub> was defined as the selectivity index (SI). The compounds of formula (I) were shown to inhibit HIV-1 effectively. Particular IC<sub>50</sub>, CC<sub>50</sub> and SI values are listed in Table 7 hereinbelow.

Table 7

Co. No.	IC <sub>50</sub> ( $\mu\text{M}$ )	CC <sub>50</sub> ( $\mu\text{M}$ )	SI
1	0.027	49.7	1860
2	0.035	>100	>2890
3	0.016	37.4	2558
4	0.315	>100	>317
5	0.094	56.2	598
6	0.020	24.4	1192
7	0.975	>100	>102
8	8.147	>100	>12
9	0.037	58.6	1587
10	2.529	>100	>39
12	1.683	55.1	32
13	0.005	7.8	1557
14	2.183	89.0	40
15	0.003	9.0	2857
16	0.389	41.2	105
17	0.167	9.1	54

Co. No.	IC <sub>50</sub> ( $\mu\text{M}$ )	CC <sub>50</sub> ( $\mu\text{M}$ )	SI
56	0.0023	1.9	839
57	0.0007	0.8	1153
58	0.0029	> 100	> 34482
59	0.0012	> 100	> 83333
60	0.29	> 100	> 350
61	0.0007	0.1	155
62	0.0032	8.7	2716
63	0.0017	0.3	198
64	0.12	> 100	> 840
65	0.18	0.2	1
66	0.0085	19.9	2347
67	0.0024	0.4	152
68	0.001	1.4	1367
69	0.0004	4.7	11632
70	0.0006	5.8	9641
71	0.0063	45.8	7275

Co. No.	IC50 ( $\mu$ M)	CC50 ( $\mu$ M)	SI
18	2.1	59.9	29
19	0.006	53.6	8642
20	0.026	36.5	1413
21	0.017	50.6	2910
22	0.035	12.2	346
23	0.001	47.9	59935
24	0.020	54.0	2667
25	0.079	>100	>1272
26	0.011	33.5	2990
27	0.017	>20	>1169
28	0.079	>20	>253
29	0.015	>20	>1324
30	0.088	>20	>228
31	0.024	86.8	3630
32	0.403	>100	>248
33	0.042	43.4	1038
34	0.319	57.8	181
35	0.103	42.3	409
36	0.323	>100	>309
37	0.443	33.4	75
38	2.449	32.4	13
39	1.531	>100	>65
40	0.253	40.2	158
41	1.986	34.2	17
42	0.352	35.5	88
43	0.603	>100	>165
45	0.010	56.3	5688
46	45.2	>100	>2
47	0.004	>100	>27027
48	44.2	>100	>1
49	0.058	45.2	786
50	0.518	52.0	100
51	0.452	>100	>221
52	0.001	2.08	2314
53	0.0006	1.3	2111

Co. No.	IC50 ( $\mu$ M)	CC50 ( $\mu$ M)	SI
72	0.0007	0.5	705
73	0.0036	> 100	> 27777
74	0.010	> 100	> 9523
75	0.0021	1.9	911
76	0.0033	5.2	1580
77	0.0030	9.6	3188
78	0.0028	0.4	144
79	0.0031	4.8	1547
80	0.011	8.7	771
81	0.0011	> 100	> 90909
82	0.0026	0.4	151
83	0.0008	0.4	541
84	0.012	9.3	753
85	0.002	0.4	208
86	0.010	> 100	> 9803
87	0.0031	2.2	711
88	0.0027	2.1	767
89	0.0007	0.4	619
90	3.4	30.8	9
91	0.0025	4.9	1976
92	45.0	> 90.0	> 2
93	0.0035	48.1	13743
94	0.0022	11.1	5064
95	0.0006	7.7	12783
96	0.0031	5.8	1885
97	0.032	13.2	415
98	2.0	13.8	7
99	0.16	59.7	367
100	0.075	0.8	10
101	0.053	29.5	558
102	0.0082	0.5	63
103	0.022	> 100	4555
104	0.0034	18.6	5476
105	52.1	< 52.1	< 1

#### D. Composition examples

The following formulations exemplify typical pharmaceutical compositions suitable for systemic administration to animal and human subjects in accordance with the present invention.

- 5 "Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) or a pharmaceutically acceptable addition salt thereof.

#### Example D.1 : film-coated tablets

##### Preparation of tablet core


- 10 A mixture of 100 g of the A.I., 570 g lactose and 200 g starch was mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole was mixed well and compressed into tablets,  
15 giving 10.000 tablets, each comprising 10 mg of the active ingredient.

##### Coating

- To a solution of 10 g methyl cellulose in 75 ml of denaturated ethanol there was added a solution of 5 g of ethyl cellulose in 150 ml of dichloromethane. Then there were added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene  
20 glycol was molten and dissolved in 75 ml of dichloromethane. The latter solution was added to the former and then there were added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated color suspension and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Claims.

1. The use of a compound of formula



- 5 a *N*-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein  
 A is CH, CR<sup>4</sup> or N;  
 n is 0, 1, 2, 3 or 4;  
 Q is hydrogen or -NR<sup>1</sup>R<sup>2</sup>;
- 10 R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl wherein each of the  
 15 aforementioned C<sub>1-12</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonylamino, mono- or di(C<sub>1-6</sub>alkyl)amino, aryl and Het; or  
 R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)aminoC<sub>1-4</sub>alkylidene;
- 20 R<sup>3</sup> is hydrogen, aryl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-6</sub>alkyloxycarbonyl; and  
 each R<sup>4</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, amino-carbonyl, nitro, amino, trihalomethyl, trihalomethyloxy or C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;
- 25 R<sup>5</sup> is hydrogen or C<sub>1-4</sub>alkyl;  
 L is C<sub>1-10</sub>alkyl, C<sub>3-10</sub>alkenyl, C<sub>3-10</sub>alkynyl, C<sub>3-7</sub>cycloalkyl, or C<sub>1-10</sub>alkyl substituted with one or two substituents independently selected from C<sub>3-7</sub>cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two, three, four or where possible five substituents each independently selected from halo,  
 30 hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, C<sub>1-6</sub>alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl, trihalomethyloxy and C<sub>1-6</sub>alkylcarbonyl; or  
 L is -X<sup>1</sup>-R<sup>6</sup> or -X<sup>2</sup>-Alk-R<sup>7</sup> wherein

$R^6$  and  $R^7$  each independently are phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyloxycarbonyl, formyl, cyano, aminocarbonyl, nitro, amino, trihalomethyloxy and trihalomethyl; and

5  $X^1$  and  $X^2$  are each independently  $-NR^3-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-S-$ ,  $-S(=O)-$  or  $-S(=O)_2-$ ;

Alk is  $C_{1-4}$ alkanediyl;

aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, nitro and

10 trifluoromethyl;


Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic hetero-

15 cyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy;


for the manufacture of a medicine for the treatment of subjects suffering from HIV (Human Immunodeficiency Virus) infection.

20 2. The use of a compound as claimed in claim 1 wherein  $n$  is at least 1 and at least one  $R^4$  is cyano.

3. The use of a compound as claimed in claim 1 or 2 wherein the compound has the formula




25 4. The use of a compound as claimed in any one of claims 1 to 3 wherein the compound has the formula



5. A compound of formula

-42-



a *N*-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein L, Q, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and A are as defined under formula (I), and R<sup>4'</sup> is halo, C<sub>1-6</sub>alkyl, cyano, aminocarbonyl, nitro, trihalomethyl, trihalomethoxy or


5 C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;

n' is 0, 1, 2 or 3;

with the proviso that Q and L are other than anilino, 2,4,6-trinitro-anilino, 3-methoxy-anilino, 4-methoxy-anilino, 3,4-dimethoxy-anilino, 3-chloro-4-fluoro-anilino, 4-cyano-anilino, 2-(C<sub>1-6</sub>alkyl)-anilino, 4-(C<sub>1-6</sub>alkyl)-anilino, 3-chloro-anilino, 4-bromo-anilino,

10 4-nitro-anilino and 4-chloro-anilino.

6. A compound as claimed in claim 5 wherein the compound has the formula



7. A compound as claimed in claim 6 wherein L and Q are other than anilino, 2,4,6-trinitro-anilino, 4-(C<sub>1-6</sub>alkyl)-anilino, 4-bromo-anilino, 4-nitro-anilino and

15 4-chloro-anilino.

8. A compound as claimed in claim 6 or 7 wherein R<sup>4'</sup> is cyano, aminocarbonyl or C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl.

9. A compound as claimed in any one of claims 5 to 8 wherein L is 2,6-dichlorobenzyl, or L is -X<sup>1</sup>-R<sup>6</sup> wherein X<sup>1</sup> is -NR<sup>3</sup>-, -S- or -O- and R<sup>6</sup> is 2,4,6-trichlorophenyl, 2,4,6-trimethyl-phenyl, 2,4-dibromo-3,5-dichloro-phenyl, 2,4-dibromo-6-fluoro-phenyl, 2,4-dichloro-6-methyl-phenyl, 2,6-dibromo-4-isopropyl-phenyl, 2,6-dibromo-4-methylphenyl, 2,6-dibromo-4-prop-1-yl-phenyl, 2,6-dichloro-4-cyano-phenyl, 2,6-dichloro-4-trifluoromethoxy-phenyl, 2,6-dichloro-4-trifluoromethyl-phenyl, 2,6-dichloro-phenyl, 2,6-dimethyl-4-(1,1-dimethylethyl)-phenyl,


20 2,6-dimethyl-phenyl, 2-bromo-4-fluoro-6-methyl-phenyl, 2-bromo-6-chloro-4-fluoro-phenyl, 4-bromo-2,6-dimethyl-phenyl, 4-chloro-2,6-dimethyl-phenyl, 4-cyano-2,6-dimethyl-phenyl; or L is -X<sup>2</sup>-Alk-R<sup>7</sup> wherein -X<sup>2</sup>-Alk- is -NH-CH<sub>2</sub>- and R<sup>7</sup> phenyl.

25

10. A compound as claimed in any one of claims 5 to 9 wherein Q is hydrogen, L is -X<sup>1</sup>-R<sup>6</sup> wherein X<sup>1</sup> is -NH- and R<sup>6</sup> is 2,4,6-trimethyl-phenyl or 4-cyano-2,6-dimethylphenyl, the NR<sup>3</sup> (optionally substituted phenyl or pyridyl) moiety represents *p*-cyano-anilino and is in the 2 position of the pyrimidine ring.
- 5 11. A compound as claimed in any one of claims 5 to 10 wherein R<sup>4'</sup> is halo, cyano, aminocarbonyl or cyanoC<sub>1-6</sub>alkyl; n is zero, A is CH, R<sup>3</sup> is hydrogen; R<sup>5</sup> is hydrogen or methyl; Q is hydrogen or NHR<sup>1</sup>; and L contains phenyl, 2,4,6-trichloro-phenyl, 2,4,6-trimethyl-phenyl, 2,4-dibromo-3,5-dichloro-phenyl, 2,4-dibromo-6-fluoro-phenyl, 2,4-dichloro-6-methyl-phenyl, 2,6-dibromo-4-isopropyl-phenyl, 10 2,6-dibromo-4-methyl-phenyl, 2,6-dibromo-4-prop-1-yl-phenyl, 2,6-dichloro-4-cyano-phenyl, 2,6-dichloro-4-trifluoromethoxy-phenyl, 2,6-dichloro-4-trifluoromethyl-phenyl, 2,6-dichloro-phenyl, 2,6-dimethyl-4-(1,1-dimethylethyl)-phenyl, 2,6-dimethyl-phenyl, 2-bromo-4-fluoro-6-methyl-phenyl, 2-bromo-6-chloro-4-fluoro-phenyl, 4-bromo-2,6-dimethyl-phenyl, 4-chloro-2,6-dimethyl-phenyl, or 15 4-cyano-2,6-dimethyl-phenyl.
12. A compound as claimed in claim 5 wherein the compound is  
 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]benzonitrile;  
 6-[(2,6-dichlorophenyl)methyl]-N2-(4-fluorophenyl)-2,4-pyrimidinediamine;  
 4-[[4-[(2,4-dichlorophenyl)methyl]-6-[(4-hydroxybutyl)amino]-2-pyrimidinyl]amino]-  
 20 benzonitrile;  
 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[(3-hydroxypropyl)amino]-2-pyrimidinyl]amino]-benzonitrile;  
 N-[2-[(4-cyanophenyl)amino]-6-[(2,6-dichlorophenyl)methyl]-4-pyrimidinyl]-  
 acetamide;  
 25 N-[2-[(4-cyanophenyl)amino]-6-[(2,6-dichlorophenyl)methyl]-4-pyrimidinyl]-  
 butanamide;  
 4-[[2-amino-6-(2,6-dichlorophenoxy)-4-pyrimidinyl]amino]benzonitrile;  
 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[(2-hydroxy-2-phenylethyl)amino]-2-  
 pyrimidinyl]amino]benzonitrile;  
 30 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[[3-(2-oxo-1-pyrrolidinyl)propyl]amino]-2-  
 pyrimidinyl]amino]benzonitrile;  
 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[[2-(2-hydroxyethoxy)ethyl]amino]-2-  
 pyrimidinyl]amino]benzonitrile;  
 35 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[(2,3-dihydroxypropyl)amino]-2-pyrimidinyl]-  
 amino]benzonitrile;


- 4-[[4-[(2,6-dichlorophenyl)methyl]-6-(hydroxyamino)-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[4-[(2-cyanoethyl)amino]-6-[(2,6-dichlorophenyl)methyl]-2-pyrimidinyl]amino]-benzonitrile;
- 5 4-[[4-[(2,6-dichlorophenyl)methyl]-6-[[2-(1-pyrrolidinyl)ethyl]amino]-2-pyrimidinyl]-amino]benzonitrile;
- 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-5-methyl-2-pyrimidinyl]amino]-benzonitrile;
- N2-(4-bromophenyl)-6-[(2,6-dichlorophenyl)methyl]-5-methyl-2,4-pyrimidinediamine;
- 10 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[2-[(2,4,6-trimethylphenyl)amino]-4-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-(2,4,6-trimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,6-dichlorophenyl)thio]-2-pyrimidinyl]amino]benzonitrile;
- 15 4-[[4-[[2,6-dibromo-4-(1-methylethyl)phenyl]amino]-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[4-[[2,6-dichloro-4-(trifluoromethyl)phenyl]amino]-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[4-[(2,4-dichloro-6-methylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 20 4-[[2-[(cyanophenyl)amino]-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;
- 4-[[4-[(2,4-dibromo-6-fluorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-5-methyl-2-pyrimidinyl]amino]-benzeneacetonitrile;
- 4-[[4-[methyl(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 25 4-[[4-[(2,4,6-trichlorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile ;
- 4-[[4-[(2,4,6-trimethylphenyl)thio]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-amino-6-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[2-amino-6-[(2,4,6-trimethylphenyl)amino]-4-pyrimidinyl]amino]benzonitrile;
- 30 4-[[4-(2-bromo-4-chloro-6-methylphenoxy)-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(4-chloro-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 3,5-dichloro-4-[[2-[(4-cyanophenyl)amino]-4-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[[2,6-dichloro-4-(trifluoromethoxy)phenyl]amino]-2-pyrimidinyl]amino]-benzonitrile;
- 35 4-[[4-[(2,4-dibromo-3,6-dichlorophenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,6-dibromo-4-propylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzamide;

- 4-[[4-[(4-(1,1-dimethylethyl)-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[2-[(4-cyanophenyl)amino]-4-pyrimidinyl]oxy]-3,5-dimethylbenzonitrile;
- 4-[[4-[(4-chloro-2,6-dimethylphenyl)amino]-5-methyl-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;
- 4-[[4-[[4-(1,1-dimethylethyl)-2,6-dimethylphenyl]amino]-5-methyl-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(4-bromo-2,6-dimethylphenyl)amino]-5-methyl-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[5-methyl-4-[(2,4,6-trimethylphenyl)thio]-2-pyrimidinyl]amino]benzonitrile;
- 4-[[4-[(2,6-dibromo-4-propylphenyl)amino]-5-methyl-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzamide, *N*3-oxide; *N*2-(4-chlorophenyl)-*N*4-(2,4,6-trimethylphenyl)-2,4-pyrimidinediamine;
- 4-[[4-[[2,6-dibromo-4-(1-methylethyl)phenyl]amino]-5-methyl-2-pyrimidinyl]amino]-benzonitrile;
- 4-[[2-[(4-cyanophenyl)amino]-5-methyl-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;
- 4-[[4-[(phenylmethyl)amino]-2-pyrimidinyl]amino]benzonitrile;
- a *N*-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof.
13. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically active amount of a compound as defined in any one of claims 5 to 12.
14. A process for preparing a pharmaceutical composition as defined in claim 13 characterized in that a therapeutically effective amount of a compound as defined in any one of claims 5 to 12 is intimately mixed with a pharmaceutically acceptable carrier.
15. A process for preparing a compound as defined in claim 5, characterized by
- a) reacting an intermediate of formula (II-A) wherein  $W^1$  is a suitable leaving group with an amino derivative of formula (III) optionally in a solvent under a reaction-inert atmosphere, and optionally in the presence of an acid




wherein Q, R<sup>3</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>5</sup>, A, n' and L are as defined in claim 5;

b) reacting an intermediate of formula (II-B) wherein W<sup>1</sup> is a suitable leaving group with an amino derivative of formula (VI) optionally in a solvent under a reaction-inert atmosphere, and optionally in the presence of an acid




c) reacting an intermediate H-X<sup>1</sup>-R<sup>6</sup> with an intermediate of formula (II-C) in a suitable solvent optionally in the presence of a suitable acid or base; thus obtaining compounds of formula (I') wherein L is -X<sup>1</sup>-R<sup>6</sup>, said compounds being represented by formula (I'-c).



or if desired, converting compounds of formula (I') into each other following art-known transformations, and further, if desired, converting the compounds of formula (I'), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms or *N*-oxides thereof..

16. A compound as claimed in claim 5 for use as a medicine.

17. The combination of a compound of formula (I) as defined in claim 1 and another antiretroviral compound.
18. The combination of a compound of formula (I') as defined in claim 5 and another antiretroviral compound.
19. A combination as claimed in claim 17 or 18 for use as a medicine.
20. A product containing (a) a compound of formula (I) as defined in claim 1, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in anti-HIV treatment.
21. A product containing (a) a compound of formula (I') as defined in claim 5, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in anti-HIV treatment.
22. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound of formula (I) as defined in claim 1, and (b) another antiretroviral compound.
23. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound of formula (I') as defined in claim 5, and (b) another antiretroviral compound.
24. A compound of formula (II'-B)



wherein  $R^3$ ,  $R^4$ ,  $R^{4'}$ ,  $n'$ ,  $A$  and  $L$  are as defined in claim 5 and  $W^1$  is a halogen.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/02043

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D239/50 C07D239/48 C07D239/46 C07D239/42 C07D401/12  
A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 91 18887 A (SMITH KLINE) 12 December 1991 (1991-12-12) cited in the application claims 1-3; examples 1-38 ---	1-4, 13-23
Y	EP 0 588 762 A (CIBA-GEIGY) 23 March 1994 (1994-03-23) page 3, line 39 - line 42; claims ---	1-4, 13-23
A	WO 95 10506 A (DU PONT MERCK) 20 April 1995 (1995-04-20) page 1 - page 13; claims ---	1-4, 13-23
X	claims; examples 4,7,32; table 1 ---	5,6,13, 16,24
A	EP 0 270 111 A (KUMIAI) 8 June 1988 (1988-06-08) the whole document ---	5,24
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

14 July 1999

Date of mailing of the international search report

29/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/02043

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DOLLY GOSH:            "2,4-BIS(ARYLAMINO)-5-METHYLPYRIMIDINES AS            ANTIMICROBIAL AGENTS."            JOURNAL OF MEDICINAL CHEMISTRY.,            vol. 10, September 1969 (1969-09), pages            974-975, XP002109182            AMERICAN CHEMICAL SOCIETY. WASHINGTON., US            ISSN: 0022-2623            page 974 - page 975</p>	<p>1-6,            13-23</p>
A	<p>CHEN CHEN ET AL.: "DESIGN AND SYNTHESIS            OF A SERIE OF NON-PEPTIDE HIGH AFFINITY            HUMAN CORTICOTROPIN-RELEASING FACTOR"            JOURNAL OF MEDICINAL CHEMISTRY.,            vol. 39, 1996, pages 4358-4360,            XP002109183            AMERICAN CHEMICAL SOCIETY. WASHINGTON., US            ISSN: 0022-2623            page 4358 - page 4359</p>	<p>1-6,            13-23</p>
A	<p>CHEMICAL ABSTRACTS, vol. 95, no. 11,            1981            Columbus, Ohio, US;            abstract no. 97712f,            D.GOSH:            "2,4-BIS(ARYLAMINO)-6-METHYLPYRIMIDINES AS            ANTIMICROBIAL AGENTS."            page 648;            XP002109184            abstract            &amp; J.INDIAN CHEM. SOC.,            vol. 58, no. 5, 1981, pages 512-513,            INDIA</p>	<p>1-6,            13-23</p>

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/02043

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9118887	A	12-12-1991	AU 7971691 A	31-12-1991
EP 588762	A	23-03-1994	JP 6184116 A	05-07-1994
			US 5516775 A	14-05-1996
WO 9510506	A	20-04-1995	AU 692484 B	11-06-1998
			AU 8012294 A	04-05-1995
			BR 9407799 A	06-05-1997
			CA 2174080 A	20-04-1995
			CN 1142817 A	12-02-1997
			CZ 9601014 A	13-11-1996
			EP 0723533 A	31-07-1996
			FI 961599 A	07-06-1996
			HR 940664 A	31-12-1996
			HU 74464 A	30-12-1996
			JP 9504520 T	06-05-1997
			NO 961425 A	12-06-1996
			NZ 274978 A	27-04-1998
			PL 313973 A	05-08-1996
			SK 47096 A	01-10-1996
			ZA 9407921 A	11-04-1996
EP 270111	A	08-06-1988	JP 2065338 C	24-06-1996
			JP 7084445 B	13-09-1995
			JP 63141971 A	14-06-1988
			US 4992438 A	12-02-1991